

- 1, p. 148, Apr. 2017, doi: 10.1186/s13104-017-2471-6.
- [10] K. Markošová, L. Weignerová, M. Rosenberg, V. Křen, and M. Rebroš, "Upscale of recombinant α -L-rhamnosidase production by *Pichia pastoris* Mut(S) strain," *Front. Microbiol.*, vol. 6, p. 1140, Oct. 2015, doi: 10.3389/fmicb.2015.01140.
- [11] V. Looser, D. Lüthy, M. Straumann, K. Hecht, K. Melzoch, and K. Kovar, "Effects of glycerol supply and specific growth rate on methanol-free production of CALB by *P. pastoris*: functional characterisation of a novel promoter," *Appl. Microbiol. Biotechnol.*, vol. 101, no. 8, pp. 3163–3176, Apr. 2017, doi: 10.1007/s00253-017-8123-x.
- [12] X. Shi, T. Karkut, M. Chamankhah, M. Alting-Mees, S. M. Hemmingsen, and D. Hegedus, "Optimal conditions for the expression of a single-chain antibody (scFv) gene in *Pichia pastoris*," *Protein Expr. Purif.*, vol. 28, no. 2, pp. 321–330, Apr. 2003, doi: 10.1016/S1046-5928(02)00706-4.
- [13] A. A. Prabhu, V. D. Veeranki, and S. J. Dsilva, "Improving the production of human interferon gamma (hIFN- γ) in *Pichia pastoris* cell factory: An approach of cell level," *Process Biochem.*, vol. 51, no. 6, pp. 709–718, 2016, doi: <https://doi.org/10.1016/j.procbio.2016.02.007>.
- [14] M. Anggiani, I. Helianti, and A. Abinawanto, "Optimization of methanol induction for expression of synthetic gene *Thermomyces lanuginosus* lipase in *Pichia pastoris*," in *AIP Conference Proceedings*, Oct. 2018, vol. 2023, no. 1, p. 20170, doi: 10.1063/1.5064154.
- [15] L. Jia, Z. Shi, J. Yan, J. Ding, and Q. Hua, "Enhancing heterologous proteins production by MutS *Pichia pastoris* via periodic methanol induction control," *AIChE J.*, vol. 66, no. 1, p. e16798, Jan. 2020, doi: 10.1002/aic.16798.
- [16] C. Dietzsch, O. Spadiut, and C. Herwig, "A fast approach to determine a fed batch feeding profile for recombinant *Pichia pastoris* strains," *Microb. Cell Fact.*, vol. 10, Oct. 2011, doi: 10.1186/1475-2859-10-85.
- [17] S. Katla, N. Mohan, S. S. Pavan, U. Pal, and S. Sivaprakasam, "Control of specific growth rate for the enhanced production of human interferon α 2b in glycoengineered *Pichia pastoris*: process analytical technology guided approach," *J. Chem. Technol. Biotechnol.*, vol. 94, no. 10, pp. 3111–3123, Oct. 2019, doi: 10.1002/jctb.6118.
- [18] S. Polez *et al.*, "A Simplified and Efficient Process for Insulin Production in *Pichia pastoris*," *PLoS One*, vol. 11, no. 12, p. e0167207, 2016, doi: 10.1371/journal.pone.0167207.
- [19] W.-C. Liu, S. Inwood, T. Gong, A. Sharma, L.-Y. Yu, and P. Zhu, "Fed-batch high-cell-density fermentation strategies for *Pichia pastoris* growth and production," *Crit. Rev. Biotechnol.*, vol. 39, no. 2, pp. 258–271, Mar. 2019, doi: 10.1080/07388551.2018.1554620.
- [20] J. Zhang, Y. Zhao, M. Li, and T. Liu, "Optimization of defined medium for recombinant *Komagataella phaffii* expressing cyclodextrin glycosyltransferase," *Biotechnol. Prog.*, vol. 35, no. 5, p. e2867, Sep. 2019, doi: 10.1002/btpr.2867.
- [21] A. A. Prabhu, B. Mandal, and V. V. Dasu, "Medium optimization for high yield production of extracellular human interferon- γ from *Pichia pastoris*: A statistical optimization and neural network-based approach," *Korean J. Chem. Eng.*, vol. 34, no. 4, pp. 1109–1121, Apr. 2017, doi: 10.1007/s11814-016-0358-1.
- [22] D. Nurdiani, Hariyatun, N. Utami, E. Wahyu Putro, and W. Kusharyoto, "Selecting *Pichia pastoris* recombinant clones for higher secretion of human insulin precursor into the culture supernatant," in *IOP Conference Series: Earth and Environmental Science*, Feb. 2020, vol. 439, no. 1, p. 12017, doi: 10.1088/1755-1315/439/1/012017.
- [23] D. Nurdiani, Hariyatun, and W. Kusharyoto, "Secretory expression of human insulin precursor in *Pichia pastoris* employing truncated alpha-factor leader sequence and a short C-peptide," *Indones. J. Biotechnol.*, vol. 23, no. 2, pp. 102–108, 2018, doi: 10.22146/ijbiotech.38958.
- [24] G. Gallo-Oller, R. Ordoñez, and J. Dotor, "A new background subtraction method for Western blot densitometry band quantification through image analysis software," *Journal of Immunological Methods*, vol. 457, Elsevier B.V., pp. 1–5, Jun. 01, 2018, doi: 10.1016/j.jim.2018.03.004.
- [25] A. Türkanoglu Özçelik, S. Yılmaz, and M. Inan, "Pichia pastoris promoters," in *Methods in Molecular Biology*, vol. 1923, Humana Press Inc., 2019, pp. 97–112.
- [26] X. Li *et al.*, "Transcriptional analysis of impacts of glycerol transporter 1 on methanol and glycerol metabolism in *Pichia pastoris*," *FEMS Yeast Res.*, vol. 18, no. 1, p. 81, Feb. 2018, doi: 10.1093/femsyr/fox081.
- [27] M. Karbalaei, S. A. Rezaee, and H. Farsiani, "Pichia pastoris: A highly successful expression system for optimal synthesis of heterologous proteins," *Journal of Cellular Physiology*, vol. 235, no. 9, Wiley-Liss Inc., pp. 5867–5881, Sep. 01, 2020, doi: 10.1002/jcp.29583.
- [28] C. B. Matthews, A. Kuo, K. R. Love, and J. C. Love, "Development of a general defined medium for *Pichia pastoris*," *Biotechnol. Bioeng.*, vol. 115, no. 1, pp. 103–113, Jan. 2018, doi: 10.1002/bit.26440.
- [29] A. Pekarisky, S. Mihalyi, M. Weiss, A. Limbeck, and O. Spadiut, "Depletion of boric acid and cobalt from cultivation media: Impact on recombinant protein production with *Komagataella phaffii*," *Bioengineering*, vol. 7, no. 4, pp. 1–24, Dec. 2020, doi: 10.3390/bioengineering7040161.
- [30] W. Zhu, R. Xu, G. Gong, L. Xu, Y. Hu, and L. Xie, "Medium optimization for high yield production of human serum albumin in *Pichia pastoris* and its efficient purification," *Protein Expr. Purif.*, vol. 181, p. 105831, May 2021, doi: 10.1016/j.pep.2021.105831.
- [31] A. B. Zepeda, A. Pessoa, and J. G. Farias, "Carbon metabolism influenced by promoters and temperature used in the heterologous protein production using *Pichia pastoris* yeast," *Brazilian J. Microbiol.*, vol. 49, pp. 119–127, Nov. 2018, doi: 10.1016/j.bjm.2018.03.010.
- [32] X. Garcia-Ortega, N. Adelantado, P. Ferrer, J. L. Montesinos, and F. Valero, "A step forward to improve recombinant protein production in *Pichia pastoris*: From specific growth rate effect on protein secretion to carbon-starving conditions as advanced strategy," *Process Biochem.*, vol. 51, no. 6, pp. 681–691, 2016, doi: <https://doi.org/10.1016/j.procbio.2016.02.018>.
- [33] C. Canales, C. Altamirano, and J. Berrios, "The growth of *Pichia pastoris* Mut+ on methanol-glycerol mixtures fits to interactive dual-limited kinetics: model development and application to optimised fed-batch operation for heterologous protein production," *Bioprocess Biosyst. Eng.*, vol. 41, no. 12, pp. 1827–1838, Dec. 2018, doi: 10.1007/s00449-018-2005-1.
- [34] D. A. Peña, B. Gasser, J. Zanghellini, M. G. Steiger, and D. Mattanovich, "Metabolic engineering of *Pichia pastoris*," *Metabolic Engineering*, vol. 50, Academic Press Inc., pp. 2–15, Nov. 01, 2018, doi: 10.1016/j.ymben.2018.04.017.
- [35] L. D. Picotto, G. H. Sguazza, M. A. Tizzano, C. M. Galosi, S. F. Cavalitto, and M. R. Pecoraro, "An effective and simplified DO-stat control strategy for production of rabies glycoprotein in *Pichia pastoris*," *Protein Expr. Purif.*, vol. 132, pp. 124–130, Apr. 2017, doi: 10.1016/j.pep.2017.02.004.