

- morphological and anatomical characters," *Biodiversitas*, vol. 20, no. 8, pp. 2340–2347, 2019, doi: 10.13057/biodiv/d200832.
- [29] U. B. Husnudin, Suharyanto, B. S. Daryono, and P. Purnomo, "Variation and non-formal classification of Indonesian eggplant (*Solanum melongena* L.) accessions based on macro and micro-morphological characters," *Agrivita*, vol. 41, no. 3, pp. 544–560, 2019, doi: 10.17503/agrivita.v41i3.2039.
- [30] C. Luo, X. H. He, H. Chen, S. J. Ou, and M. P. Gao, "Analysis of diversity and relationships among mango cultivars using Start Codon Targeted (SCoT) markers," *Biochem. Syst. Ecol.*, vol. 38, no. 6, pp. 1176–1184, 2010, doi: 10.1016/j.bse.2010.11.004.
- [31] E. Zulaika, L. Sembiring, and A. Soegianto, "Characterization and Identification Of Mercury-resistant Bacteria From Kalimas River Surabaya-Indonesia By Numerical Phenetic Taxonomy," *J. Basic. Appl. Sci. Res.*, vol. 2, no. 7, pp. 7263–7269, 2012, [Online]. Available: www.textroad.com.
- [32] M. K. Fleshman *et al.*, "Carotene and novel apocarotenoid concentrations in orange-fleshed cucumis melo melons: Determinations of β -carotene bioaccessibility and bioavailability," *J. Agric. Food Chem.*, vol. 59, no. 9, pp. 4448–4454, 2011, doi: 10.1021/jf200416a.
- [33] B. S. Daryono, S. D. Maryanto, S. Nissa, and G. R. Aristya, "Analisis Kandungan Vitamin Pada Melon (*Cucumis melo* L.) Kultivar Melodi Gama 1 dan Melon Komersial," *Biog. J. Ilm. Biol.*, vol. 4, no. 1, pp. 1–9, 2016, doi: 10.24252/bio.v4i1.1113.