

REFERENCES

- [1] Motley, T. J., The ethnobotany of Fagraea Thunb. (Gentianaceae) : The timber of Malesia and the scent of Polynesia, *Economic Botany* 2004, 58 (3), 396-409.
- [2] Backlund M, Oxelman B, Bremer B: Phylogenetic relationships within the Gentianales based on ndhF and rbcL sequences, with particular reference to the Loganiaceae: *american journal of botany* 2000, 87(7):1029–1043.
- [3] Kristiati, A. N., 2008, Buku Ajar Fitokimia, Airlangga University Press: Surabaya
- [4] Ibrahim, S., 1998, Teknik Laboratorium Kimia Organik, Pasca Sarjana Universitas Andalas. Hal. 19, 35-42.
- [5] Kumar, H. V. K., Navyashree, S. N., Rakshitha, H. R. dan Chauhan, J. B., 2012, Studies on the free radical scavenging activity of Syagrus romanzoffiana, *International journal of pharmaceutical and biomedical research*, Vol. 3 (2), 81-84
- [6] Bhatt Mehl K, Dholwani Kishor K and Saluja Ajay K, Isolation and structure elucidation of Scopoletin from *Ipomoea reniformis* (Convolvulaceae), *Journal of Applied Pharmaceutical Science* 01 (05); 2011: 138-144.
- [7] Murray RDH, Mendez J, Brown SA: *The Natural Coumarins*. Occurrence, Chemistry and Biochemistry, John Wiley and Sons, United Kingdom, 1982, 28 – 31.