













- [32] S. Gujral, K. P. C. Sharma, P. Gupta, and A. Wani, "Anti-oxidants properties of legumes and their morphological fractions as affected by cooking," *Food Sci. Biotechnol.*, vol. 22, no. 1, pp. 187–194, 2013.
- [33] D. F. Holderbaum, T. Kon, and M. P. Guerra, "Scientia Horticulturae Dynamics of total phenolic content in different apple tissues and genotypes : impacts and relevance for breeding programs," *Sci. Hortic. (Amsterdam)*, vol. 168, pp. 58–63, 2014.
- [34] S. Boukhanouf, H. Louaileche, and D. Perrin, "Phytochemical content and in vitro anti-oxidant activity of faba bean (*Vicia faba* L.) as affected by maturity stage and cooking practice," *Int. Food Res. J.*, vol. 23, no. 3, pp. 954–961, 2016.
- [35] V. E. Nambi, R. K. G. Sunil, and K. P. C. Sharma, "Degradation Kinetics of Bioactive Components, Anti-oxidant Activity, Colour and Textural Properties of Selected Vegetables During Blanching," *J. Food Sci. Technol.*, vol. 53, no. 7, pp. 3073–3082, 2016.
- [36] A. Nirbaya, F. R. Zakaria, and H. Wijaya, "Stability of Bioactive Compound And Fatty Acid Composition After Thermal Processing of Winged Bean (*Psophocarpus tetragonolobus*)," *Int. J. Adv. Sci. Technol.*, vol. 29, no. 7, pp. 1602–1610, 2020.
- [37] R. M. A. Khalili *et al.*, "Total Phenolic Content and In Vitro Antioxidant Activity of Winged Bean (*Psophocarpus tetragonolobus*)," *Pakistan J. Nutr.*, vol. 12, no. 5, pp. 416–422, 2013.
- [38] N. Nazir, M. Zahoor, and M. Nisar, "A Review on Traditional Uses and Pharmacological Importance of Genus *Elaeagnus* Species," *Bot. Rev.*, vol. 86, no. 3–4, pp. 247–280, Dec. 2020.
- [39] D. Hou *et al.*, "Mung Bean (*Vigna radiata* L.): Bioactive Polyphenols, Polysaccharides, Peptides, and Health Benefits," *Nutrients*, vol. 11, no. 6, p. 1238, May 2019.