

REFERENCES

- [1] G.R. Gibson and M.B. Roberfroid, Dietary modulation of the human colonic microbiota: Introducing the concept of prebiotics. *The Journal of nutrition*, vol. 125 (6), pp. 1401-1412, 1995.
- [2] D.C. Montrose and M.H. Floch, Probiotics used in human studies. *Journal of clinical gastroenterology*, vol. 39(6), pp. 469-484, 2005.
- [3] R. Sahadeva, S. Leong, K. Chua, C. Tan, et al., Survival of commercial probiotic strains to pH and bile. *International Food Research Journal*, vol. 18(4), 2011.
- [4] L. Siew-Wai, T. Zi-Ni, A.A. Karim, N.M. Hani, et al., Fermentation of Metroxylon sagu resistant starch type III by *Lactobacillus* sp. and *Bifidobacterium bifidum*. *Journal Of Agricultural and Food Chemistry*, vol. 58(4), pp. 2274-2278, 2010.
- [5] N. Hussain, M.A. Othaman, and M.K. Ayob, Effects of Inulin and Inulin Containing Chocolate on in vitro Growth of Bifidobacteria. *Sains Malaysiana*, vol. 41(10), pp. 1223-1231, 2012.
- [6] A.M. Boehm, Tailored drapery tab. 2005, Google Patents.
- [7] H. Chung and D.L. Kasper, Microbiota-stimulated immune mechanisms to maintain gut homeostasis. *Current opinion in immunology*, vol. 22(4), pp. 455-460, 2010.
- [8] S. Ding, M.M. Chi, B.P. Scull, R. Rigby, et al., High-fat diet: bacteria interactions promote intestinal inflammation which precedes and correlates with obesity and insulin resistance in mouse. *PLoS one*, vol. 5(8), 2010.
- [9] F. Bäckhed, J.K. Manchester, C.F. Semenkovich and J.I. Gordon, Mechanisms underlying the resistance to diet-induced obesity in germ-free mice. *Proceedings of the National Academy of Sciences*, vol. 104(3), pp. 979-984, 2007.
- [10] R. Havenaar, Intestinal health functions of colonic microbial metabolites: A review. *Beneficial microbes*, vol. (2), pp. 103-114, 2011.
- [11] P.D. Cani, A.M. Neyrinck, F. Fava, C. Knaut, et al., Selective increases of bifidobacteria in gut microflora improve high-fat-diet-induced diabetes in mice through a mechanism associated with endotoxaemia. *Diabetologia*, vol. 50(11), pp. 2374-2383, 2007.
- [12] M. Hill, Intestinal flora and endogenous vitamin synthesis. *European journal of cancer prevention: the official journal of the European Cancer Prevention Organisation (ECP)*, vol. 6, pp. S43-5, 1997.
- [13] K.E. Scholz-Ahrens, P. Ade, B. Marten, P. Weber, et al., Prebiotics, probiotics, and synbiotics affect mineral absorption, bone mineral content, and bone structure. *The Journal of nutrition*, vol. 137(3), pp. 838S-846S, 2007.
- [14] P. Markowiak and K. Śliżewska, Effects of probiotics, prebiotics, and synbiotics on human health. *Nutrients*, vol. 9(9), pp. 1021, 2017.
- [15] T.C. Wallace, F. Guarner, K. Madsen, M.D. Cabana, et al., Human gut microbiota and its relationship to health and disease. *Nutrition reviews*, vol. 69(7), pp. 392-403, 2012.
- [16] S. Bogdanov, Honey as nutrient and functional food. *Proteins*, vol. 1100, pp. 1400-2700, 2012.
- [17] S.J. Mei, M.M. Nordin, and N.A. Sani, Fructooligosaccharides in honey and effects of honey on growth of *Bifidobacterium longum* BB 536. *International Food Research Journal*, vol. 17(3), pp. 557-561, 2010.
- [18] L.P. Oddo, M. Piazza, A. Sabatini, and M. Accorti, Characterization of unifloral honeys. *Apidologie*, vol. 26(6), pp. 453-465, 1995.
- [19] R.J. Weston and L.K. Brocklebank, The oligosaccharide composition of some New Zealand honeys. *Food Chemistry*, vol. 64(1), pp. 33-37, 1999.
- [20] Z. Ustunol and H. Gandhi, Growth and viability of commercial *Bifidobacterium* spp. in honey-sweetened skim milk. *Journal of Food Protection*, vol. 64(11), pp. 1775-1779, 2001.
- [21] J.H. Cummings, G.T. Macfarlane, and H.N. Englyst, Prebiotic digestion and fermentation. *The American journal of clinical nutrition*, vol. 73(2), pp. 415-420, 2001.
- [22] K. Bemmo, M. Sahoo, R. Jayabalan, and N. Zambou, Honey, probiotics and prebiotics: review. *Res J Pharm Biol Chem Sci*, 2016. 7: p. 2428-2438.
- [23] L.M. Tsai, Studies on *Acacia mangium* in Kemasul forest, Malaysia. I. Biomass and productivity. *Journal of Tropical Ecology*, 1988: p. 293-302.
- [24] P. Höglberg, and J. Wester, Root biomass and symbioses in *Acacia mangium* replacing tropical forest after logging. *Forest ecology and management*, 1998. 102(2-3): p. 333-338.
- [25] A. Galiana, et al., Nitrogen fixation estimated by the 15N natural abundance method in *Acacia mangium* Willd. inoculated with *Bradyrhizobium* sp. and grown in silvicultural conditions. *Soil Biology and Biochemistry*, 2002. 34(2): p. 251-262.
- [26] V. Morales, M. Sanz, A. Olano, and N. Corzo, Rapid separation on activated charcoal of high oligosaccharides in honey. *Chromatographia*, vol. 64(3-4), pp. 1-6, 2006.
- [27] M.L. Sanz, N. Polemis, V. Morales, N. Corzo, et al., In vitro investigation into the potential prebiotic activity of honey oligosaccharides. *Journal Of Agricultural and Food Chemistry*, vol. 53(8), pp. 2914-2921, 2005.
- [28] L. Curda and M. Plockova, Impedance measurement of growth of lactic acid bacteria in dairy cultures with honey addition. *International Dairy Journal*, vol. 5(7), pp. 727-733, 1995.
- [29] A. Ting and J. Decosta, Comparison of the viability of probiotics from various cultured-milk drinks in a simulated pH study of the human gastrointestinal tract. *International Food Research Journal*, vol. 16(1), pp. 59-64, 2009.
- [30] M. Moniruzzaman, M.I. Khalil, S.A. Sulaiman, and S.H. Gan, Physicochemical and antioxidant properties of Malaysian honeys produced by *Apis cerana*, *Apis dorsata* and *Apis mellifera*. *BMC Complementary and Alternative Medicine*, vol. 13(1), pp. 43, 2013.
- [31] W. Scheppach, Effects of short chain fatty acids on gut morphology and function. *Gut*, vol.35, pp. S35-S38, 1994.