











- [20] K. Sanatombi and S. Rajkumari, "Effect of Processing on Quality of Pepper: A Review," *Food Rev. Int.*, vol. 36, no. 6, pp. 626–643, 2020, doi: 10.1080/87559129.2019.1669161.
- [21] K. H. Murti, "The effect of drying temperature on vitamin C content of Lado F1 curly chili fruit (*Capsicum annuum* L)," *J. Keteknikan Pertan. Trop. dan Biosist.*, vol. 5, no. 3, pp. 245–256, 2017.
- [22] J. A. V. Costa, B. C. B. Freitas, L. Moraes, M. Zapparoli, and M. G. Morais, "Progress in the physicochemical treatment of microalgae biomass for value-added product recovery," *Bioresour. Technol.*, vol. 301, p. 122727, 2020, doi: 10.1016/j.biortech.2019.122727.
- [23] Y. P. Wanita, Purwaningsih, and M. Tri, "Value-added diversification of red chili processing into dried chili," in *Prosiding Seminar Nasional Hasil Penelitian Pangan dan Hasil Pertanian 2018*, 2018, pp. 480–489, [Online]. Available: <https://semnas.tp.ugm.ac.id/prosiding/>.
- [24] C. Nimrotham, R. Songprakorp, S. Thepa, and V. Monyakul, "Experimental research of drying red chili by two methods: solar drying and low-temperature system drying," *Energy Procedia*, vol. 138, pp. 512–517, 2017, doi: 10.1016/j.egypro.2017.10.237.
- [25] T. Naemsai, J. Jareanjit, and K. Thongkaew, "Experimental investigation of solar-assisted heat pump dryer with heat recovery for the drying of chili peppers," *J. Food Process Eng.*, vol. 42, no. 6, pp. 1–10, 2019, doi: 10.1111/jfpe.13193.
- [26] K. Rybak, A. Wiktor, D. Witrowa-Rajchert, O. Parniakov, and M. Nowacka, "The effect of traditional and non-Thermal treatments on the bioactive compounds and sugars," *Molecules*, vol. 25, pp. 1–17, 2020.
- [27] L. . Montoya-Ballesteros, A. Gonzales-Leon, Y. . Martinez-Nunez, M. . Robles-Burgueno, M. . Garcia-Alvarado, and G. . Rodriguez-Jimenes, "Impact of open sun-drying and hot air drying on capsaicin, capsanthin, and ascorbic acid content in chiltepin (*capsicum annum* L. var. *glabriusculum*)," *Rev. Mex. Ing. Quimica*, vol. 16, no. 3, pp. 813–825, 2017, [Online]. Available: <http://www.redalyc.org/articulo.oa?id=62029966013>.
- [28] N. Azizzuddin and A. Abdullah, "Effect of blanching treatments on antioxidant activity of frozen green capsicum (*Capsicum annum* L. var bell pepper)," *Int. Food Res. J.*, vol. 25, no. 4, pp. 1427–1434, 2018, doi: 10.1063/1.4966775.
- [29] I. O. Nyangena, W. O. Owino, S. Imathiu, and J. Ambuko, "Effect of pretreatments prior to drying on antioxidant properties of dried mango slices," *Sci. African*, vol. 6, p. e00148, 2019, doi: 10.1016/j.sciaf.2019.e00148.
- [30] V. K. Maurya, K. M. Gothandam, V. Ranjan, A. Shakya, and S. Pareek, "Effect of drying methods (microwave-vacuum, freeze, hot air, and sun-drying) on physical, chemical and nutritional attributes of five pepper (*Capsicum annum* var. *annuum*) cultivars," *J. Sci. Food Agric.*, vol. 98, no. 9, pp. 3492–3500, 2018, doi: 10.1002/j.
- [31] Indonesian National Standardization Agency, *SNI 7388:2009 Maximum limit of microbial contamination in food*. 2009.