

authors are obliged to the research team that supervise the data analysis using chemometrics tools and the e-nose operation.

REFERENCES

- [1] C. Viet, M. Nguyet, T. Nguyen, M. Hanh, T. Truong, and X. Dong, "Screening for Optimal Parameters of Nattokinase Synthesis by *Bacillus subtilis* natto in Solid-State Fermentation," vol. 10, no. 3, pp. 1207–1213, 2020.
- [2] S. L. Percival and D. W. Williams, *Escherichia coli*, Second Edi. Elsevier, 2013.
- [3] J. A. Crump and J. Wain, *Salmonella*, Third Edit. Elsevier Inc., 2016.
- [4] F. Fung, H. S. Wang, and S. Menon, "Food safety in the 21st century," *Biomed. J.*, vol. 41, no. 2, pp. 88–95, 2018, doi: 10.1016/j.bj.2018.03.003.
- [5] M. K. Sharif, K. Javed, and A. Nasir, *Foodborne Illness: Threats and Control*, vol. 15. Elsevier Inc., 2018.
- [6] M. Falasconi, I. Concina, E. Gobbi, V. Sberveglieri, A. Pulvirenti, and G. Sberveglieri, "Electronic Nose for Microbiological Quality Control of Food Products," *Int. J. Electrochem.*, vol. 2012, pp. 1–12, 2012, doi: 10.1155/2012/715763.
- [7] J. M. Cevallos-Cevallos, M. D. Danyluk, and J. I. Reyes-De-Corcuera, "GC-MS Based Metabolomics for Rapid Simultaneous Detection of *Escherichia coli* O157:H7, *Salmonella typhimurium*, *Salmonella muenchen*, and *Salmonella hartford* in Ground Beef and Chicken," *J. Food Sci.*, vol. 76, no. 4, pp. 238–246, 2011, doi: 10.1111/j.1750-3841.2011.02132.x.
- [8] Y. X. Yu and Y. Zhao, "Electronic Nose Integrated with Chemometrics for Rapid Identification of Foodborne Pathogen," *Chemom. Pract. Appl.*, 2012, doi: 10.5772/32099.
- [9] E. Tait, J. D. Perry, S. P. Stanforth, and J. R. Dean, "Identification of Volatile Organic Compounds Produced by Bacteria Using HS-SPME-GC-MS," *J. Chromatogr. Sci.*, vol. 52, no. 4, pp. 363–373, 2014, doi: 10.1093/chromsci/bmt042.
- [10] A. P. F. Turner and N. Magan., "Electronic Noses and Disease Diagnostics: Nature Reviews," *Microbiol. (Nature Rev.)*, vol. 2, no. February, pp. 1–7, 2004, doi: 10.1038/nrmicro823.
- [11] X. Tian, J. Wang, and S. Cui, "Analysis of Pork Adulteration in Minced Mutton Using Electronic Nose of Metal Oxide Sensors," *J. Food Eng.*, vol. 119, no. 4, pp. 744–749, 2013, doi: 10.1016/j.jfoodeng.2013.07.004.
- [12] J. Carrillo and C. Durán, "Fast identification of Bacteria for Quality Control of Drinking Water through A Static Headspace Sampler Coupled to a Sensory Perception System," *Biosensors*, vol. 9, no. 1, p. 23, 2019, doi: 10.3390/bios9010023.
- [13] M. R. Cavallari *et al.*, "A hybrid electronic nose and tongue for the detection of ketones: Improved sensor orthogonality using graphene oxide-based detectors," no. CVD, 2017, doi: 10.1109/JSEN.2017.2661067.
- [14] J. Fitzgerald and H. Fenniri, "Cutting edge methods for non-invasive disease diagnosis using e-tongue and e-nose devices," *Biosensors*, vol. 7, no. 4, 2017, doi: 10.3390/bios7040059.
- [15] J. A. Covington, E. W. Westenbrink, N. Ouaret, R. Harbord, K. D. Bardhan, and R. P. Arasaradnam, "Application of a Novel Tool for Diagnosing Bile Acid Diarrhoea," pp. 11899–11912, 2013, doi: 10.3390/s130911899.
- [16] P. F. Astantri, W. S. A. Prakoso, K. Triyana, T. Untari, C. M. Airin, and P. Astuti, "Lab-made electronic nose for fast detection of *Listeria monocytogenes* and *Bacillus cereus*," *Vet. Sci.*, vol. 7, no. 1, pp. 1–11, 2020, doi: 10.3390/vetsci7010020.
- [17] U. Siripatrawan, "Rapid Differentiation between *E. coli* and *Salmonella Typhimurium* Using Metal Oxide Sensors Integrated with Pattern Recognition," *Sensors Actuators, B Chem.*, vol. 133, no. 2, pp. 414–419, 2008, doi: 10.1016/j.snb.2008.02.046.
- [18] G. C. Green, A. D. C. Chan, H. Dan, and M. Lin, "Using a metal oxide sensor (MOS)-based electronic nose for discrimination of bacteria based on individual colonies in suspension," *Sensors Actuators, B Chem.*, vol. 152, no. 1, pp. 21–28, 2011, doi: 10.1016/j.snb.2010.09.062.
- [19] S. Balasubramanian, S. Panigrahi, C. M. Logue, H. Gu, and M. Marchello, "Neural networks-integrated metal oxide-based artificial olfactory system for meat spoilage identification," *J. Food Eng.*, vol. 91, no. 1, pp. 91–98, 2009, doi: 10.1016/j.jfoodeng.2008.08.008.
- [20] I. A. Casalnuovo, D. Di Pierro, M. Coletta, and P. Di Francesco, "Application of Electronic Noses for Disease Diagnosis and Food Spoilage Detection," pp. 1428–1439, 2006.
- [21] J. Yan *et al.*, "Electronic nose feature extraction methods: A review," *Sensors (Switzerland)*, vol. 15, no. 11, pp. 27804–27831, 2015, doi: 10.3390/s151127804.
- [22] S. N. Hidayat *et al.*, "The Electronic Nose Coupled with Chemometric Tools for Discriminating the Quality of Black Tea Samples In Situ," *Chemosensors*, vol. 7, no. 3, p. 29, Jul. 2019, doi: 10.3390/chemosensors7030029.
- [23] S. N. Hidayat and K. Triyana, "Optimized Back-Propagation Combined with Radial Basic Neural Network for Improving Performance of the Electronic Nose: Case Study on the Fermentation Process of Tempeh," vol. 1755, pp. 7–11, 2016, doi: 10.1063/1.4958466.
- [24] K. Triyana, M. Tauhid Subekti, P. Aji, S. Nur Hidayat, and A. Rohman, "Development of Electronic Nose with Low-Cost Dynamic Headspace for Classifying Vegetable Oils and Animal Fats," *Appl. Mech. Mater.*, vol. 771, pp. 50–54, 2015, doi: 10.4028/www.scientific.net/amm.771.50.
- [25] S. Panigrahi, S. Balasubramanian, H. Gu, C. Logue, and M. Marchello, "Neural-Network-Integrated Electronic Nose System for Identification of Spoiled Beef," *LWT - Food Sci. Technol.*, vol. 39, no. 2, pp. 135–145, 2006, doi: 10.1016/j.lwt.2005.01.002.
- [26] H. Elgaali, T. R. Hamilton-Kemp, M. C. Newman, R. W. Collins, K. Yu, and D. D. Archbold, "Comparison of long-chain alcohols and other volatile compounds emitted from food-borne and related Gram positive and Gram negative bacteria," *J. Basic Microbiol.*, vol. 42, no. 6, pp. 373–380, 2002, doi: 10.1002/1521-4028(200212)42:6<373::AID-JOBM373>3.0.CO;2-4.
- [27] N. Karami *et al.*, "Initial study of three different pathogenic microorganisms by gas chromatography-mass spectrometry," *F1000Research*, vol. 6, no. 0, p. 1415, 2018, doi: 10.12688/f1000research.12003.3.
- [28] U. Siripatrawan and B. R. Harte, "Solid phase microextraction/gas chromatography/mass spectrometry integrated with chemometrics for detection of *Salmonella typhimurium* contamination in a packaged fresh vegetable," *Anal. Chim. Acta*, vol. 581, no. 1, pp. 63–70, 2007, doi: 10.1016/j.aca.2006.08.007.
- [29] K. P. Talaro and B. Chess, *Foundations in Microbiology*, 8th ed. New York: McGraw-Hill Companies, Inc, 2012.
- [30] H. H. Nguyen, S. Y. Yi, A. Woubit, and M. Kim, "A Portable Surface Plasmon Resonance Biosensor for Rapid Detection of *Salmonella typhimurium*," *Appl. Sci. Conver. Technol.*, vol. 25, no. 3, pp. 61–65, 2016, doi: 10.5757/asct.2016.25.3.61.
- [31] A. T. Arfao, C. L. Djimeli, O. V. N. Ewoti, M. Nola, and T. Sime-ngando, "Detachment of adhered enteropathogenic *Escherichia coli* cells from polythene fragments immersed in aquatic microcosm using *Eucalyptus microcorys* extract (Myrtaceae)," no. May 2016, 2017.
- [32] I. Tazi, K. Triyana, D. Siswanta, A. C. A. Veloso, A. M. Peres, and L. G. Dias, "Dairy Products Discrimination According to the Milk Type Using an Electrochemical Multisensor Device Coupled with Chemometric Tools," *J. Food Meas. Charact.*, vol. 12, no. 4, pp. 2385–2393, 2018, doi: 10.1007/s11694-018-9855-8.