















- hepatopancreatic necrosis disease,” *Aquac. Reports*, vol. 18, 2020, doi: 10.1016/j.aqrep.2020.100442.
- [19] B. V. Chang *et al.*, “Investigation of a farm-scale multitrophic recirculating aquaculture system with the addition of *Rhodovulum sulfidophilum* for milkfish (*Chanos chanos*) coastal aquaculture,” *Sustain.*, vol. 11, no. 7, 2019, doi: 10.3390/su11071880.
- [20] A. Omont, R. Elizondo-González, E. Quiroz-Guzmán, C. Escobedo-Fregoso, R. Hernández-Herrera, and A. Peña-Rodríguez, “Digestive microbiota of shrimp *Penaeus vannamei* and oyster *Crassostrea gigas* co-cultured in integrated multi-trophic aquaculture system,” *Aquaculture*, vol. 521, p. 735059, 2020, doi: 10.1016/j.aquaculture.2020.735059.
- [21] D. Kaya, E. Genc, M. A. Genc, M. Aktas, O. T. Eroldogan, and D. Guroy, “Biofloc technology in recirculating aquaculture system as a culture model for green tiger shrimp, *Penaeus semisulcatus*: Effects of different feeding rates and stocking densities,” *Aquaculture*, vol. 528, p. 735526, 2020, doi: 10.1016/j.aquaculture.2020.735526.
- [22] F. G. Addo, S. Zhang, B. Manirakiza, O. E. Ohore, and Y. Shudong, “The impacts of straw substrate on biofloc formation, bacterial community and nutrient removal in shrimp ponds,” *Bioresour. Technol.*, vol. 326, no. November 2020, p. 124727, 2021, doi: 10.1016/j.biortech.2021.124727.
- [23] T. N. D. Khoa, C. T. Tao, L. Van Khanh, and T. N. Hai, “Super-intensive culture of white leg shrimp (*Litopenaeus vannamei*) in outdoor biofloc systems with different sunlight exposure levels: Emphasis on commercial applications,” *Aquaculture*, vol. 524, p. 735277, 2020, doi: 10.1016/j.aquaculture.2020.735277.
- [24] Z. Chen *et al.*, “Effects of carbon source addition on microbial community and water quality in recirculating aquaculture systems for *Litopenaeus vannamei*,” *fish. Sci.*, vol. 86, no. 3, pp. 507–517, 2020, doi: 10.1007/s12562-020-01423-3.
- [25] L. S. Herbeck, U. Krumme, I. Nordhaus, and T. C. Jennerjahn, “Pond aquaculture effluents feed an anthropogenic nitrogen loop in a SE Asian estuary,” *Sci. Total Environ.*, vol. 756, p. 144083, 2021, doi: 10.1016/j.scitotenv.2020.144083.
- [26] W. Xu *et al.*, “Effects of feeding frequency on growth, feed utilization, digestive enzyme activity and body composition of *Litopenaeus vannamei* in biofloc-based zero-exchange intensive systems,” *Aquaculture*, vol. 522, no. January, 2020, doi: 10.1016/j.aquaculture.2020.735079.
- [27] H. Lemonnier and S. Faninoz, “Effect of water exchange on effluent and sediment characteristics and on partial nitrogen budget in semi-intensive shrimp ponds in New Caledonia,” *Aquac. Res.*, vol. 37, no. 9, pp. 938–948, 2006, doi: 10.1111/j.1365-2109.2006.01515.x.
- [28] M. Morshed, S. Islam, H. Das Lohano, and P. Shyamsundar, “Production externalities of shrimp aquaculture on paddy farming in coastal Bangladesh,” *Agric. Water Manag.*, vol. 238, no. May, p. 106213, 2020, doi: 10.1016/j.agwat.2020.106213.
- [29] J. G. Cardoso-Mohedano, F. Páez-Osuna, F. Amezcua-Martínez, A. C. Ruiz-Fernández, G. Ramírez-Reséndiz, and J. A. Sanchez-Cabeza, “Combined environmental stress from shrimp farm and dredging releases in a subtropical coastal lagoon (SE Gulf of California),” *Mar. Pollut. Bull.*, vol. 104, no. 1–2, pp. 83–91, 2016, doi: 10.1016/j.marpolbul.2016.02.008.
- [30] N. T. M. Pratiwi, B. Widigdo, and D. A. Syifa, “Water quality and organic content from intensive system of vaname production at coastal area of Sumur, Pandeglang, Banten,” *IOP Conf. Ser. Earth Environ. Sci.*, vol. 420, no. 1, 2020, doi: 10.1088/1755-1315/420/1/012022.
- [31] M. R. Islam and S. Tabeta, “Shrimp vs prawn-rice farming in Bangladesh: A comparative impacts study on local environments and livelihoods,” *Ocean Coast. Manag.*, vol. 168, no. November 2018, pp. 167–176, 2019, doi: 10.1016/j.ocecoaman.2018.11.004.
- [32] V. Hukom, R. Nielsen, M. Asmild, and M. Nielsen, “Do Aquaculture Farmers Have an Incentive to Maintain Good Water Quality? The Case of Small-Scale Shrimp Farming in Indonesia,” *Ecol. Econ.*, vol. 176, no. May, p. 106717, 2020, doi: 10.1016/j.ecolecon.2020.106717.
- [33] O. M. Joffre, J. R. De Vries, L. Klerkx, and P. M. Poortvliet, “Why are cluster farmers adopting more aquaculture technologies and practices? The role of trust and interaction within shrimp farmers’ networks in the Mekong Delta, Vietnam,” *Aquaculture*, vol. 523, no. October 2019, p. 735181, 2020, doi: 10.1016/j.aquaculture.2020.735181.
- [34] K. C. Tran, A. Dalsgaard, P. T. Van, and B. P. Tersbøl, “To pray in four directions: Understanding Vietnamese farmers’ shrimp health management practices,” *Aquaculture*, vol. 536, no. February 2020, 2021, doi: 10.1016/j.aquaculture.2021.736406.
- [35] U. Senarath and C. Visvanathan, “Environmental issues in brackish water shrimp aquaculture in Sri Lanka,” *Environ. Manage.*, vol. 27, no. 3, pp. 335–348, 2001, doi: 10.1007/s002670010153.