











- [13] K. C. Chinsembu, "Diabetes mellitus and nature's pharmacy of putative antidiabetic plants," *J. Herb. Med.*, vol. 15, p. 100230, Mar. 2019.
- [14] P. Somwong and C. Chuchote, "Determination of Lupeol, a Cytotoxic Compound Against SW620 Cells in the Extracts of Ha-Rak Recipe," *Pharmacogn. J.*, vol. 13, no. 1, pp. 133–141, 2021.
- [15] J. Lohakul *et al.*, "The protective effect of polyherbal formulation, Harak Formula, on UVA-induced photoaging of human dermal fibroblasts and mouse skin via promoting Nrf2-regulated antioxidant defense," *Front. Pharmacol.*, vol. 12, p. 649820, Apr. 2021.
- [16] C. N. Johnson *et al.*, "Biodiversity losses and conservation responses in the Anthropocene," *Science (80- )*, vol. 356, no. 6335, pp. 270–275, Apr. 2017.
- [17] W. J. Ripple, C. Wolf, T. M. Newsome, M. Hoffmann, A. J. Wirsing, and D. J. McCauley, "Extinction risk is most acute for the world's largest and smallest vertebrates," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 114, no. 40, pp. 10678–10683, Oct. 2017.
- [18] A. Qin *et al.*, "Maxent modeling for predicting impacts of climate change on the potential distribution of *Thuja sutchuenensis* Franch., an extremely endangered conifer from southwestern China," *Glob. Ecol. Conserv.*, vol. 10, pp. 139–146, Apr. 2017.
- [19] J. C. Deb, S. Phinn, N. Butt, and C. A. McAlpine, "The impact of climate change on the distribution of two threatened Dipterocarp trees," *Ecol. Evol.*, vol. 7, no. 7, pp. 2238–2248, Apr. 2017.
- [20] A. Norberg *et al.*, "A comprehensive evaluation of predictive performance of 33 species distribution models at species and community levels," *Ecol. Monogr.*, vol. 89, no. 3, p. 1370, Aug. 2019.
- [21] E. Kaky, V. Nolan, A. Alatawi, and F. Gilbert, "A comparison between Ensemble and MaxEnt species distribution modelling approaches for conservation: A case study with Egyptian medicinal plants," *Ecol. Inform.*, vol. 60, p. 101150, Nov. 2020.
- [22] M. Abdelaal, M. Fois, G. Fenu, and G. Bacchetta, "Using MaxEnt modeling to predict the potential distribution of the endemic plant *Rosa arabica* Crép. in Egypt," *Ecol. Inform.*, vol. 50, pp. 68–75, Mar. 2019.
- [23] K. Zhang *et al.*, "Impact of climate factors on future distributions of *Paeonia ostii* across China estimated by MaxEnt," *Ecol. Inform.*, vol. 50, pp. 62–67, Mar. 2019.
- [24] N. Xu, F. Meng, G. Zhou, Y. Li, B. Wang, and H. Lu, "Assessing the suitable cultivation areas for *Scutellaria baicalensis* in China using the Maxent model and multiple linear regression," *Biochem. Syst. Ecol.*, vol. 90, p. 104052, Jun. 2020.
- [25] L. Marshall *et al.*, "The interplay of climate and land use change affects the distribution of EU bumblebees," *Glob. Chang. Biol.*, vol. 24, no. 1, pp. 101–116, Jan. 2018.
- [26] M. S. Hosseinzadeh, F. Q. Masood, N. Babak, R. Dennis, and S. M. Kazemi, "Habitat suitability and modelling the potential distribution of the Plateau Snake Skink *Ophiomorus nuchalis* (Sauria scincidae) on the Iranian Plateau," *North. West. J. Zool.*, vol. 14, no. 1, pp. 60–63, 2018.
- [27] A. Yudaputra, I. P. Astuti, and W. P. Cropper, "Comparing six different species distribution models with several subsets of environmental variables: Predicting the potential current distribution of *Guettarda speciosa* in Indonesia," *Biodiversitas J. Biol. Divers.*, vol. 20, no. 8, pp. 2321–2328, Jul. 2019.
- [28] A. Yudaputra, "Modelling potential current distribution and future dispersal of an invasive species *Calliandra calothyrsus* in Bali Island, Indonesia," *Biodiversitas J. Biol. Divers.*, vol. 21, no. 2, pp. 674–682, Jan. 2020.
- [29] D. A. Pisner and D. M. Schnyer, "Support vector machine," in *Machine Learning: Methods and Applications to Brain Disorders*, Elsevier, 2019, pp. 101–121.
- [30] J. M. Kass, S. I. Meenan, N. Tinoco, S. F. Burneo, and R. P. Anderson, "Improving area of occupancy estimates for parapatric species using distribution models and support vector machines," *Ecol. Appl.*, vol. 31, no. 1, p. e02228, Jan. 2021.
- [31] A. Yudaputra, I. Fijridiyanto, and W. P. Cropper, "The potential impact of climate change on the distribution pattern of *Eusideroxylon zwageri* (Bornean Ironwood) in Kalimantan, Indonesia," *Biodiversitas*, vol. 21, no. 1, pp. 326–333, 2020.
- [32] F. Ghareghan, G. Ghanbarian, H. R. Pourghasemi, and R. Safaeian, "Prediction of habitat suitability of *Morina persica* L. species using artificial intelligence techniques," *Ecol. Indic.*, vol. 112, p. 106096, May 2020.
- [33] O. Ovaskainen *et al.*, "How to make more out of community data? A conceptual framework and its implementation as models and software," *Ecol. Lett.*, vol. 20, no. 5, pp. 561–576, May 2017.
- [34] F. K. C. Hui, "Boral: Bayesian ordination and regression analysis," *R package version 1.4*, 2017. [Online]. Available: <https://cran.r-project.org/package=boral>. [Accessed: 25-May-2021].
- [35] A. Gayen and H. R. Pourghasemi, "Spatial modeling of gully erosion: A new ensemble of CART and GLM data-mining algorithms," in *Spatial Modeling in GIS and R for Earth and Environmental Sciences*, Elsevier, 2019, pp. 653–669.
- [36] T. Connor *et al.*, "Effects of grain size and niche breadth on species distribution modeling," *Ecography (Cop.)*, vol. 41, no. 8, pp. 1270–1282, Aug. 2018.
- [37] T. H. Booth, H. A. Nix, J. R. Busby, and M. F. Hutchinson, "Bioclim: The first species distribution modelling package, its early applications and relevance to most current MaxEnt studies," *Divers. Distrib.*, vol. 20, no. 1, pp. 1–9, Jan. 2014.
- [38] S. E. Fick and R. J. Hijmans, "WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas," *Int. J. Climatol.*, vol. 37, no. 12, pp. 4302–4315, Oct. 2017.
- [39] R. J. Hijmans, S. Phillips, J. Leathwick, and J. Elith, "Dismo: Species distribution modeling," *R Package Version 1.3-3*, 2020. [Online]. Available: <https://cran.r-project.org/web/packages/dismo/index.html>. [Accessed: 24-May-2021].
- [40] J. C. Cooper and J. Soberón, "Creating individual accessible area hypotheses improves stacked species distribution model performance," *Glob. Ecol. Biogeogr.*, vol. 27, no. 1, pp. 156–165, Jan. 2018.
- [41] W. J. Krzanowski and D. J. Hand, *ROC: Curves for continuous data*. Boca Raton, FL: Chapman and Hall/CRC, 2009.
- [42] T. Handayani, "Diversity, potential and conservation of Annonaceae in Bogor Botanic Gardens, Indonesia," *Biodiversitas*, vol. 19, no. 2, pp. 591–603, 2018.
- [43] J. E. Teysmann, *Catalogus plantarum quae in Horto Botanico Bogoriensi coluntur*. Batavia: Lands-Drukkerij, 1866.
- [44] D. Setyanti and D. O. Pribadi, "Mapping the dynamics of dead trees collection to support sustainable landscape management at Bogor Botanic Gardens," in *IOP Conference Series: Earth and Environmental Science*, 2020, pp. 1–6.
- [45] D. A. Lestari, R. Azrianingsih, and H. Hendrian, "Taxonomical position of Annonaceae species from East Java, Indonesia: Collections of Purwodadi botanic garden based on morphological character," *Biodiversitas*, vol. 18, no. 3, pp. 1067–1076, Jul. 2017.
- [46] D. A. Lestari, A. P. Fiqa, and I. K. Abywijaya, "Leaf morphological traits of *Orophea* spp. (Annonaceae): Living collections of Purwodadi Botanic Gardens, East Java, Indonesia," *Biodiversitas J. Biol. Divers.*, vol. 22, no. 6, pp. 3403–3411, Jun. 2021.
- [47] T. Handayani, Yuzammi, and J. T. Hadiah, "Inflorescence morphology and development of Suweg (*Amorphophallus paeoniifolius* (Dennst.) Nicolson)," *Biodiversitas*, vol. 21, no. 12, pp. 5835–5844, Dec. 2020.
- [48] F. Damayanti, R. Vitri Garvita, H. Wawangningrum, and S. Rahayu, "Flower development, pollen viability and pollen storage test of *Aeschynanthus radicans*," *Biodiversitas*, vol. 22, no. 4, pp. 1940–1945, Apr. 2021.
- [49] D. A. Lestari and A. P. Fiqa, "Environmental factors influence on flowering and fruiting period of selected essential oil plants from annonaceae," *Biodiversitas*, vol. 21, no. 3, pp. 910–921, Mar. 2020.
- [50] B. Moungsrimuangdee, P. Waiboonya, P. Larpkern, P. Yods-Nga, and M. Saeyang, "Reproductive phenology and growth of riparian species along Phra Prong river, Sa Kaeo province, Eastern Thailand," *J. Landsc. Ecol.*, vol. 10, no. 2, pp. 35–48, 2017.