















- study,” *J. Crop Improv.*, vol. 32, no. 5, pp. 657–680, 2018, doi: 10.1080/15427528.2018.1487354.
- [31] R. Huang, Y. Wang, K. Li, and Y. Q. Wang, “Genetic variation and population structure of clonal Zingiber zerumbet at a fine geographic scale: a comparison with two closely related selfing and outcrossing Zingiber species,” *BMC Ecol. Evol.*, vol. 21, no. 1, pp. 1–13, 2021, doi: 10.1186/s12862-021-01853-2.
- [32] Wahyuni, S., D. H. Xu, N. Bermawie, H. Tsunematsu, and T. Ban., “Genetic relationships among ginger accessions based on AFLP marker,” *J. Bioteknol. Pertan.*, vol. 8, no. 2, pp. 60–68, 2003.
- [33] V. Kumar and B. K. Roy, “Population authentication of the traditional medicinal plant Cassia tora L. based on ISSR markers and FTIR analysis,” *Sci. Rep.*, vol. 8, no. 1, pp. 1–11, 2018, doi: 10.1038/s41598-018-29114-1.
- [34] K. Ashraf, S. Sultan, and S. A. A. Shah, “Phychemistry, Phytochemical, Pharmacological and Molecular Study of Zingiber Officinale Roscoe: a Review,” *Int. J. Pharm. Pharm. Sci.*, vol. 9, no. 10, p. 8, 2017, doi: 10.22159/ijpps.2017v9i11.19613.
- [35] A. District, “Genetic diversity encourages conservation of threatened ethno medicinally important plant species in Koz Abakhel District Swat , KP , Pakistan,” no. July, pp. 35–46, 2019.
- [36] H. Purnomo, F. Jaya, and S. B. Widjanarko, “The effects of type and time of thermal processing on ginger (*Zingiber officinale* Roscoe) rhizome antioxidant compounds and its quality,” *Int. Food Res. J.*, vol. 17, no. 2, pp. 335–347, 2010.
- [37] S. Diehn, B. Zimmermann, V. Tafintseva, M. Bağcıoğlu, A. Kohler, M. Ohlson, S. Fjellheim, and J. Kneipp, “Discrimination of grass pollen of different species by FTIR spectroscopy of individual pollen grains,” *Analytical and Bioanalytical Chemistry*, vol. 412, no. 24, pp. 6459–6474, Apr. 2020, doi: 10.1007/s00216-020-02628-2.
- [38] B.-J. Lee, Y. Zhou, J. S. Lee, B. K. Shin, J.-A. Seo, D. Lee, Y.-S. Kim, and H.-K. Choi, “Discrimination and prediction of the origin of Chinese and Korean soybeans using Fourier transform infrared spectrometry (FT-IR) with multivariate statistical analysis,” *PLOS ONE*, vol. 13, no. 4, p. e0196315, Apr. 2018, doi: 10.1371/journal.pone.0196315.
- [39] Y. Y. Wang, J. Q. Li, H. G. Liu, and Y. Z. Wang, “Attenuated total reflection-Fourier transform infrared spectroscopy (ATR-FTIR) combined with chemometrics methods for the classification of Lingzhi species,” *Molecules*, vol. 24, no. 12, 2019, doi: 10.3390/molecules24122210.
- [40] D. Gaikwad, S. K. Sachin, K. V. Ashwini, S. Jadhav, and M. Gadhve, “Isolation and standardization of gingerol from ginger rhizome by using TLC, HPLC, and identification tests,” *Pharma Innov. J.*, vol. 6, no. 2, pp. 179–182, 2017.
- [41] A. Ardila, I. Chairani, N. Nurdianti, and N. H. Fitriyah, “Fabrikasi Nanopartikel Herbal Dalam Tablet Effervescent Menggunakan Metode Solvent Emulsification Diffusion Kombinasi High Speed Homogenizer,” *Pros. Semnastek*, no. Prosiding Semnastek 2017, pp. 1–8, 2017.
- [42] L. Triyasmono, I. Munisa, K. Anwar, T. Wianto, and H. B. Santoso, “Identification and authentication of *Eurycoma longifolia* root extract from *Zingiber officinale* rhizome using FTIR spectroscopy and chemometrics,” *Indones. J. Chemom. Pharm. Anal.*, pp. 1–9, 2020.
- [43] R. Kooke, L. Morgado, F. Becker, H. van Eekelen, R. Hazarika, Q. Zheng, R. C. H. de Vos, F. Johannes, and J. J. B. Keurentjes, “Epigenetic mapping of the *Arabidopsis* metabolome reveals mediators of the epigenotype-phenotype map,” *Genome Research*, vol. 29, no. 1, pp. 96–106, Nov. 2018, doi: 10.1101/gr.232371.117.