

ACKNOWLEDGMENT

We thank LPPM Universitas Brawijaya for providing grants for junior researchers/ Hibah Peneliti Pemula (HPP). We also thank Aprilia Nur Komariyah and Ilham Putra Adiyaksa for helping during data validation.

REFERENCES

- [1] Y. R. Yanza *et al.*, "Coleus amboinicus (Lour.) leaves as a modulator of ruminal methanogenesis and biohydrogenation in vitro¹," *J. Anim. Sci.*, vol. 96, no. 11, pp. 4868–4881, Nov. 2018.
- [2] S. Ślusarczyk *et al.*, "Phytochemical profile and antioxidant activities of coleus amboinicus Lour. Cultivated in Indonesia and Poland," *Molecules*, vol. 26, no. 10, 2021.
- [3] R. Damanik, "Torbangun: a Batakese traditional cuisine for the lactating mother in the North Sumatra Province of Indonesia," in *Handbook of Dietary and Nutritional Aspects of Human Breast Milk*, no. 5, 2013, pp. 631–644.
- [4] R. Damanik, L. Kustiyah, M. Hanafi, and A. C. Iwansyah, "Evaluation Lactogenic Activity of Ethyl Acetate Fraction of Torbangun (Coleus amboinicus L .) Leaves," in *IOP Conference Series: Earth and Environmental Science*, 2017, pp. 2–10.
- [5] R. Damanik, "Summary Statement : Torbangun (Coleus amboinicus Lour) : A Batakese Traditional," *J Hum Lact*, vol. 25, no. 1, pp. 64–72, 2019.
- [6] T. F. Hutajulu and L. Junaidi, "The Benefit of Bangun-Bangun Leaves Extract (Coleus amboinicus L.) for Increasing of Rat Breast Milk Production," *J. Ris. Ind.*, vol. 7, no. 1, pp. 15–24, 2013.
- [7] W. Yulianto, N. Andarwulan, P. E. Giriwono, and J. Pamungkas, "Bioactive Compounds from Torbangun (Plectranthus amboinicus L. Spreng) Chloroform Fraction Induce Apoptosis in Breast Cancer (MCF-7 Cells) In Vitro," vol. 22, no. April, pp. 37–44, 2017.
- [8] N. Katurci, N. Işık, Ç. Güpür, H. O. Güler, O. Gursoy, and Y. Yılmaz, "Differences in antioxidant activity, total phenolic and flavonoid contents of commercial and homemade tomato pastes," *J. Saudi Soc. Agric. Sci.*, vol. 19, no. 4, pp. 249–254, 2020.
- [9] S. R. Dewi, N. Sumarni, N. Izza, A. W. Putranto, and B. Susilo, "Studi Variasi Kuat Medan Listrik PEF dan Metode Pengeringan Bahan Terhadap Senyawa Antioksidan Ekstrak Daun Torbangun (Coleus amboinicus L.)," *J. Keteknikan Pertanian*, vol. 7, no. 1, pp. 91–98, 2019.
- [10] S. R. Dewi, B. D. Argo, and N. Ulya, "Kandungan Flavonoid dan Aktivitas Antioksidan Ekstrak Pleurotus ostreatus," *Rona Tek. Pertanian*, vol. 11, no. 1, pp. 1–10, 2018.
- [11] Y. Zhao, Y. Li, and R. Zhang, "Silica aerogels having high flexibility and hydrophobicity prepared by sol-gel method," *Ceram. Int.*, vol. 44, no. 17, pp. 21262–21268, 2018.
- [12] F. N. Idris and M. M. Nadzir, "Comparative studies on different extraction methods of centella asiatica and extracts bioactive compounds effects on antimicrobial activities," *Antibiotics*, vol. 10, no. 4, 2021.
- [13] D. Naviglio, P. Scarano, M. Ciaravolo, and M. Gallo, "Rapid solid-liquid dynamic extraction (RSLDE): A powerful and greener alternative to the latest solid-liquid extraction techniques," *Foods*, vol. 8, no. 7, pp. 1–22, 2019.
- [14] N. Medina-Torres, T. Ayora-Talavera, H. Espinosa-Andrews, A. Sánchez-Contreras, and N. Pacheco, "Ultrasound assisted extraction for the recovery of phenolic compounds from vegetable sources," *Agronomy*, vol. 7, no. 3, 2017.
- [15] G. Xu, X. Ye, J. Chen, and D. Liu, "Effect of heat treatment on the phenolic compounds and antioxidant capacity of citrus peel extract," *J. Agric. Food Chem.*, vol. 55, no. 2, pp. 330–335, 2007.
- [16] N. Izza and D. Trisanti, "The optimization of ultrasonic-assisted extraction of antioxidant compounds from butterfly pea flower (Clitoria ternatea L.) by using response surface methodology," *IOP Conf. Ser. Earth Environ. Sci.*, vol. 743, no. 1, p. 012046, May 2021.
- [17] N. Izza, S. R. Dewi, A. W. Putranto, and D. Yulneri, "Extraction of Phenolic Compounds from Cosmos caudatus Using Pulse Electric Field (PEF)," *J. Teknol. Pertanian*, vol. 17, no. 2, pp. 91–96, 2016.
- [18] K. Ahmad Shiekh, O. Oduwaye Olatunde, B. Zhang, N. Huda, and S. Benjakul, "Pulsed electric field assisted process for extraction of bioactive compounds from custard apple (Annona squamosa) leaves," *Food Chem.*, vol. 359, p. 129976, 2021.
- [19] V. M. Pappas *et al.*, "Use of Pulsed Electric Field as a Low-Temperature and High-Performance 'Green' Extraction Technique for the Recovery of High Added Value Compounds from Olive Leaves," *Beverages*, vol. 7, no. 3, 2021.
- [20] M. Afroz Bakht *et al.*, "Ultrasound-assisted extraction of some branded tea: Optimization based on polyphenol content, antioxidant potential and thermodynamic study," *Saudi J. Biol. Sci.*, vol. 26, no. 5, pp. 1043–1052, 2018.
- [21] M. Bouafia *et al.*, "The optimization of ultrasonic-assisted extraction of Centaurea sp. antioxidative phenolic compounds using response surface methodology," *J. Appl. Res. Med. Aromat. Plants*, vol. 25, p. 100330, 2021.
- [22] N. Izza *et al.*, "Microwave-assisted extraction of phenolic compounds from Moringa oleifera seed as anti-biofouling agents in membrane processes," *MATEC Web Conf.*, vol. 204, 2018.
- [23] A. W. Putranto, S. R. Dewi, N. Izza, D. R. Yulneri, M. Y. S. Dachi, and S. H. Sumarlan, "Ekstraksi senyawa fenolik daun kenikir (Cosmos caudatus) menggunakan Microwave Assisted Extraction (MAE)," *Rona Tek. Pertanian*, vol. 11, no. 1, pp. 60–71, 2018.
- [24] N. Utama-ang, S. Sida, P. Wanachantararak, and A. Kawee-ai, "Development of edible Thai rice film fortified with ginger extract using microwave-assisted extraction for oral antimicrobial properties," *Sci. Rep.*, vol. 11, no. 1, p. 14870, 2021.
- [25] L. Azaroual *et al.*, "Optimization of the Microwave-Assisted Extraction of Simple Phenolic Compounds from Grape Skins and Seeds," *Agronomy*, vol. 11, no. 8, 2021.
- [26] H. Baltacıoğlu, C. Baltacıoğlu, I. Okur, A. Tanrıvermiş, and M. Yalç, "Optimization of microwave-assisted extraction of phenolic compounds from tomato: Characterization by FTIR and HPLC and comparison with conventional solvent extraction," *Vib. Spectrosc.*, vol. 113, p. 103204, Mar. 2021.
- [27] A. V. B. Reddy, M. Moniruzzaman, V. Madhavi, and J. Jaafar, "Chapter 8 - Recent improvements in the extraction, cleanup and quantification of bioactive flavonoids," in *Bioactive Natural Products*, vol. 66, B. T.-S. in N. P. C. Atta-ur-Rahman, Ed. Elsevier, 2020, pp. 197–223.
- [28] A. Pandey, T. Belwal, K. C. Sekar, I. D. Bhatt, and R. S. Rawal, "Optimization of ultrasonic-assisted extraction (UAE) of phenolics and antioxidant compounds from rhizomes of Rheum moorcroftianum using response surface methodology (RSM)," *Ind. Crops Prod.*, vol. 119, pp. 218–225, 2018.
- [29] X. Lin, L. Wu, X. Wang, L. Yao, and L. Wang, "Ultrasonic-assisted extraction for flavonoid compounds content and antioxidant activities of India Moringa oleifera L. leaves: Simultaneous optimization, HPLC characterization and comparison with other methods," *J. Appl. Res. Med. Aromat. Plants*, vol. 20, p. 100284, 2021.
- [30] S. Jafarinejad, "7 - Solid-Waste Management in the Petroleum Industry," S. B. T.-P. W. T. and P. C. Jafarinejad, Ed. Butterworth-Heinemann, 2017, pp. 269–345.
- [31] B. Dahmoune *et al.*, "Microwave assisted extraction of bioactive saponins from the starfish Echinaster sepositus: Optimization by response surface methodology and comparison with ultrasound and conventional solvent extraction," *Chem. Eng. Process. - Process Intensif.*, vol. 163, no. February, 2021.
- [32] E. J. Davis, E. Spadoni Andreani, and S. Karboune, "Production of Extracts Composed of Pectic Oligo/Polysaccharides and Polyphenolic Compounds from Cranberry Pomace by Microwave-Assisted Extraction Process," *Food Bioprocess Technol.*, pp. 634–649, 2021.
- [33] N. Kutlu, A. İsci, O. Sakiyan, and A. E. Yılmaz, "Extraction of Phenolic Compounds from Cornelian Cherry (Cornus mas L.) Using Microwave and Ohmic Heating Assisted Microwave Methods," *Food Bioprocess Technol.*, pp. 650–664, 2021.
- [34] A. C. Iwansyah, M. R. M. Damanik, L. Kustiyah, and M. Hanafi, "Potensi Fraksi Etil Asetat Daun Torbangun (Coleus amboinicus L.) dalam Meningkatkan Produksi Susu, Bobot Badan Tikus, dan Anak Tikus," *J. Gizi dan Pangan*, vol. 12, no. 1, pp. 61–68, 2017.
- [35] N. Andarwulan, N. D. Yuliana, E. Hasna, S. A. Aziz, and T. D. Davis, "Comparative Analysis of Three Torbangun Clones (–Plectranthus amboinicus– (Lour.) Spreng) Based on Phenotypic Characteristics and Phenolic Content," *Am. J. Plant Sci.*, vol. 05, no. 24, pp. 3673–3683, 2014.