

- [13] R. E. Mickens, "Calculation of denominator functions for non-standard finite difference schemes of differential equations satisfying a positivity condition," *Numerical Method for Partial Differential Equations*, pp. 672–691, 2007.
- [14] R. Anguelov, and J. M. Lubuma, "Non-standard finite difference method by non-local approximation," *Mathematics and Computers in Simulation*, pp. 465–475, 2003.
- [15] A. Ibijola, and A. A. Obayomi, "Derivation of new non-standard finite difference schemes for non-autonomous ordinary differential equation," *American Journal of Scientific and Industrial Research*, pp. 122–127, 2012.
- [16] A.A. Obayomi, "A new set of non-standard finite difference schemes for the solution of an equation of the type $y' = y(1 - y^2)$," *International Journal of Pure and Applied Sciences and Technology*, vol. 12(2), pp. 34–42, 2012.
- [17] A.A. Obayomi, and A. Olu, "Construction of new non-standard finite difference schemes for the solution of a free un-damped harmonic oscillator equation," *Mathematical Theory and Modeling*, vol. 3(2), pp. 91–97, 2013.
- [18] Z. Yaacob, and M. K. Hasan, "Non-standard finite difference schemes for natural convection in an inclined porous rectangular cavity.," in *IEEE Proc. of the Electrical Engineering and Informatics Conference*, 2015, pp. 665–669.
- [19] N. M. M. Yusop, and M. K. Hasan, "Development of new harmonic Euler using non-standard finite difference technique for solving stiff problems," *Jurnal Teknologi*, vol. 77 (20), pp. 19–24, 2015.
- [20] N. M. M. Yusop, M. K. Hasan, M. F. M. Amran, and M. Wook, "Investigation on non-standard scheme applied to Euler and third order Euler for initial value problems," *International Journal of Computer Applications*, vol. 155(14), pp. 7–10, 2016.
- [21] N. M. M. Yusop, M. K. Hasan, M. Wook, M. F. M. Amran, and S. R. Ahmad, "Comparison new algorithm modified euler based on harmonic-polygon approach for solving ordinary differential equation," *Journal of Telecommunication, Electronic and Computer Engineering*, vol. 9(2-11), pp. 29–32, 2017.
- [22] M. Abid, N. Abbas, H. Z. Nazir, and Z. Lin, "Enhancing the mean ratio estimators for estimating population mean using nonconventional location parameters," *Revista Colombiana de Estadística*, vol. 39(1), pp. 63–79, 2016.
- [23] C. Feng, and B. Vidakovic, "Estimation of the hurst exponent using trimean estimators on nondecimated wavelet coefficients," 2015. <https://arxiv.org/pdf/1709.08775>
- [24] A. Akbarinia, and A. Parraga, "Colour constancy beyond the classical receptive field," *IEEE Transaction on Pattern Analysis and Machine Intelligence*. 2016. doi: 10.1109/TPAMI.2017.2753239
- [25] S. Maqbool, and S. Javaid, "Variance estimation using linear combination of trimean and quartile average," *American Journal of Biological and Environmental Statistics*, vol. 3(1), pp. 5–9, 2017.
- [26] N. A. Othman, and M. K. Hasan, "New hybrid two-step method for simulating Lotka-Volterra model," *Pertanika Journal of Science and Technology*, vol. 25(S), pp. 115–124, 2017.
- [27] M. K. Hasan, and S. Mazlan, "Adapting non-standard weighted average approach for solving the Lotka-Volterra model," *Pertanika Journal of Science and Technology*, vol. 25(S), pp. 97–106, 2017.
- [28] F. Brauer, and C. C. Chavez, *Mathematical Models in Population Biology and Epidemiology*, F. Braur Ed. New York, United States: Springer-Verlag, 2001.
- [29] D.T. Dimitrov, and H.V. Kojauhrov, "Positive and elementary stable non-standard numerical methods with applications to predator-prey models," *Applied Mathematics Letters*, pp. 98–108, 2006