















- varieties from Thailand, China and Sri Lanka. *Food Chem.* 2011;124:132–40. 10.1016/j.foodchem.2010.05.115, 2011.
- [15] Shen Y, Jin L, Xiao P, Lu Y, Bao J. Total phenolics, flavonoids, antioxidant capacity in rice grain and their relations to grain color, size and weight. *J Cereal Sci.* 2009;49:106–11. 10.1016/j.jcs.2008.07.010, 2009.
- [16] Lukman A, Anggraini, D. Rahmawati, N. Suhaeni, N. (2013). 1-Preparation and Test properties have been praelatinized. *Indonesian Pharmaceutical Research Journal* 1 (2): 67-71.
- [17] Yoshida H, Tomiyama Y, Mizushima Y. Lipid components, fatty acids and triacylglycerol molecular species of black and red rices. *Food Chem* 2010;123:210–5. 10.1016/j.foodchem.2010.04.010, 2010.
- [18] Yodmanee, S., T.T. Karrila, dan P. Pakdeechanuan. Physical, Chemical and Antioxidant Properties of Pigmented Rice Grown in Southern Thailand. *International Food Research Journal*, 18 (3), 901-906, 2011.
- [19] Abdel-Aal, E. M., J.C. Young dan I. Rabalski. Anthocyanin Composition in Black, Blue, Pink, Purple, and Red Cereal Grains. *J. Agric. Food Chem.*, 54, 4696-4704, 2006.
- [20] Webb, G.P. *Dietary Supplements and Functional Foods*. United Kingdom: Blackwell Publishing, Ltd, 2006.
- [21] Sompong,R.,S.Siebenhandl-Ehn, G.Linsberger-Martin,dan E.Berghofer. (2011). Physicochemical and antioxidative properties of red and black rice varieties from Thailand, China and Sri Lanka. *Elsevier Appl. Sci. Pbl.*, 124, 132-140.
- [22] FD Goffman and CJ Bergman. Rice kernel phenolic content and its relationship with antiradical efficiency *Journal of the Science of Food and Agriculture Sci Food Agric* 84:1235–1240, 2004. Mangiri J, N. Mayulu, S.Kawengian. Overview of Nutrients in Black Rice (*Oryza sativa* L.) Kultivar Pare Ambo South Sulawesi, 2015.
- [23] N. Yilmaz and N. B Tuncel. The effect of infrared stabilisation on B vitamins, phenolics and antioxidants in rice bran. *interantional journal food science technology*, 2014.
- [24] Jeong, S. Moon, Youngkim, D. Ryulkim, S. Chunjo, K. O. Nam, dan S. C. Lee. Effect of Heat Treatment on the Antioxidant Activity of Extracts from Citrus peels. *J. Agric. Food Chem.* 52 : 3389-3393, 2004.
- [25] Gaulejac, N. S-C, C. Provost and N. Vivas. Comparative Study of Polyphenol Scavenging Activities Assessed by Different Methods. *Journal of Agricultural and Food Chemistry.* 47: 425-431, 1998.
- [26] AOAC (Association of Official Analytical Chemyst). *Official Method of Analysis of The Association of Official Analytical of Chemist.* Arlington (USA): The Association of Official Analytical Chemist, Inc, 2005.
- [27] Julkunen dan Tiitto, R. Phenolics Constituens in the Leaves of Northern Willows : Methods for the Analysis of Certain Phenolics. *J. Agric. Food Chem.* 33: 213-217, 1985.
- [28] Matthaus, B. 2002. Antioxidant Activity of Extracts Obtained from Residues of Different Oilseeds. *J. Agric. Food Chem.* 50: 3444-3452, 2010.
- [29] F. Pourmorad1, S. J. Hosseinimehr1, N. Shahabimajd. Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *African Journal of Biotechnology* Vol. 5 (11), pp. 1142-1145, 2 June 2006 Available online at <http://www.academicjournals.org/AJB> ISSN 1684–5315 © 2006 Academic Journals, 2006.
- [30] Walter M., Marchesan E. Phenolic compounds and antioxidant activity of rice. *Braz. Arch. Biol. Technol.* 2011;54:371–377. doi: 10.1590/S1516-89132011000200020. 2011.
- [31] Muntana, N., dan Prasong, S. Study on total phenolic contents and their antioxidant activities of thai whaite, red, and black rice bran extracts. *Pakistan Journal of Biological Sciences* 13(4): 170-174, 2010.
- [32] Vichapong J, Sookserm M, Srijesdaruk V, Swatsitang P, Srijaranai S. High Perfomance Liquid Chromatograich Analysis of Phenolic Compounds and Their Antioxidant Activities in Rie Varieties.LWT-Food Science Technology 43:1325-1330, 2010.
- [33] Khalekuzzaman, M., Datta, K., Oliva, N., Alam, M.F., Joarder, I. dan Datta, S.K. Stable integration, expression and inheritance of the ferritin gene in transgenic elite Indica rice cultivar BR29 with enhanced iron level in the endosperm. *Indian Journal of Biotechnology* 5: 26-31, 2006.
- [34] Sutharut, J. dan Sudarat, J. Total anthocyanin content and antioxidant activity of germinated colored rice. *International Food Research Journal* 10(1): 215-221, 2012.
- [35] L. Amagliani, J. O'Regan, A. L. Kelly. James A. O'Mahony. The composition, extraction, functionality and applications of rice proteins: A review. The composition, extraction, functionality and applications of rice proteins: A review. *Trends in food sciences and technology.* Volume 64, June 2017, Pages 1-12, 2017.
- [36] Masniati. Characteristics of Physicochemical Properties of Red Rice in Several Rice Production Centers in South Sulawesi. Universitas Hasanudin, 2013.
- [37] T. Chand and B. Savitri. Book *Vitamins*. First published: 29 April 2016 <https://doi.org/10.1002/9783527681754.ch3>, 2018.
- [38] Indrasari SD. The mineral content of superior varieties of rice and its relation to health. *Food Crop Science and Technology.* 1 (1): 88-99, 2006.
- [39] Kubo, A., Yugichi, Y., Suzuki, S. Satoh, H and Kitamura, S. The use of microbeam X-ray diffraction for the characterisation of starch crystal structure in rice mutant kurnels of waxy, amylose extender and sugary1. *Journal of Cereal Science*, 48, pp.92-97. 2008.
- [40] Shifeng Yu & Ying Ma & Lucile Menager & Da-Wen Sun. Physicochemical Properties of Starch and Flour from Different Rice Cultivars. *Food Bioprocess Technol* (2012) 5:626–637 DOI 10.1007/s11947-010-0330-8, 2010.
- [41] Iturriaga, L., Lopez, B., & Añon, M. Thermal and physico-chemical characterization of seven argentine rice flours and starches. *Food Research International*, 37, 439–447, 2004.
- [42] Singh, N., Nakaura, Y., Inouchi, N., & Nishinari, K. Fine structure, thermal and viscoelastic properties of starches separat-ed from indica rice cultivars. *Starch/Stärke*, 59, 10–20, 2007.
- [43] Kang, H.-J., Hwang, I.-K., Kim, K.-S., & Choi, H.-C. Comparative structure and physicochemical properties of II pumbyeo, a high-quality japonica rice, and its mutant, suweon *Journal of Agricultural and Food Chemistry*, 51, 6598– 6603, 2003.
- [44] Manful, J. T., Grimm, C. C., Gayin, J., & Coker, R. D. Effect of variable parboiling on crystallinity of rice samples. *Cereal Chemistry*, 85, 92–95. <https://doi.org/10.1094/CCHEM-85-1-0092>, 2008.
- [45] Mulyaningsih TR. The content of Fe and Zn in food products for agricultural products, livestock and fisheries using the k0-AANI method. *Indonesian Journal of Nuclear Science and Technology.* 10 (2): 71-80, 2009.
- [46] Yustisia, Tohari, Shiddieq Dj, Subowo. Enrichment of Iron (Fe) and Zinc (Zn) in Rice and Determining Characteristics of Efficient Rice Paddy Varieties in Soil Vertisols and Inceptisols. *Agrotrop: Journal on Agriculture Science.* 2 (1): 67-75, 2012.
- [47] Dianawati N, Sugiarto RD. Determination of Iron Levels During Rice Maturation Phase Using UV-Vis Spectrophotometer. *ITS Science and Art Journal.* 4 (2): 35-38, 2015
- [48] Fleming, D.E., Foran, K.A. Kim, J.S. and Guernsey, J.R. Portable Xray flourescence for assessing trace eleents in rice and rice products: Comparison with inductively-mass spectrometry. *Appl Radiant Isot*, 104217-23, doi:10.1016/j.apradiso.2015.07.014.
- [49] Eckhoff, B., Kannig Ber, B., Langhoff, N., Wadel, R. and Wolf, H. *Handbook of Practical X-Ray Fluorescence Analysis.* ISBN 3-540-28603-e 2006..
- [50] Kaur, J. and Kumar, A. Elemental analysis of different varieties of rice samples using XRF technique. *AIP Conference Proceedings* 1, vol.1728. 020350.2016. <https://doi.org/10.1063/1.4945401>