





days when combined with IAA and the soil medium of top soil+sand and with NAA with both soil media (Table 2). However, the cuttings planted with IAA and saw dust+sand soil media produced roots in more than 9 days, although not significantly different from the test.

TABLE I

SOURCE OF VARIATION, DEGREES OF FREEDOM AND MEAN SQUARES TAKEN FROM ANOVA FOR TIME TO APPEARANCE OF THE FIRST SHOOT, NUMBER OF SHOOT AT 9 AND 12 DAYS AFTER PLANTING (DAP)

Source of variation I/	df	Mean squares		
		Time to appearance of the first shoot	Number of shoots	
			9 DAP	12 DAP
Variety (V)	4	16.3583*** 2/	1.8057***	1.8730**
Media (M)	1	4.2667	0.3511	0.5332
Hormone (H)	1	0.0000	0.1354	0.1522
V*M	4	0.5583	0.0860	0.0486
V*H	4	1.7917	0.1902	0.2655
M*H	1	0.0667	0.0184	0.0070
V*M*H	4	10.3583**	0.6403*	1.0837*
Error	38	1.9026	0.1763	0.3321
CV%		13.75	36.40	28.30

1/ v – variety; m – soil media; and h – rooting hormone;  
2/ \* significant at p=0.05; \*\* : significant at p=0.01; \*\*\* : significant at p=0.001

TABLE II

THREE-WAY INTERACTION AMONG VARIETY, SOIL MEDIA AND HORMONE FOR TIME TO APPEARANCE OF THE FIRST SHOOT OF CUTTINGS OF MALAY APPLE

Variety	Hormone			
	IAA 1000 ppm		NAA 1000 ppm	
	Top soil+ Sand	Sawdust + Sand	Top soil+ Sand	Sawdust + Sand
Malay Apple Jati	9.7±1.16	12.7±0.58	13.0±1.73	11.3±1.53
Malay Apple Bottle	11.3±0.58	10.0±1.73	9.0±1.00	11.7±1.53
Malay Apple Bagea	8.0±0.0	10.0±3.46	8.0±0.0	8.0±0.0
Malay Apple Dark Purple	9.3±0.58	9.3±1.16	10.0±0.0	9.7±0.58
Malay Apple Red Circle	10.7±2.31	9.3±0.58	8.7±1.16	11.0±1.73

The malay variety Jati had significantly earlier sprouting in 9 days with IAA combined with top soil+sand medium, where as other combinations took more than 11 days. The variety Bottle ranged from 9 days in NAA combined with top soil + sand to 11 days with IAA and top soil + sand as well as with IAA and NAA with saw dust and sand mixtures. Although insignificant, Dark purple and Red Circle varieties too took 9 to 11 days for the appearance of first shoot.

### B. Number of Shoots Produced

The number of shoots produced was also influenced by a significant three-way interaction among the variety, hormone and soil media (Table 1). The number of shoots produced at 9 DAP was nearly 1 (Table 3), while, it took 12 days for the second shoot, and the number of shoots was 2 for all the varieties (Table 4). This indicates that there was no difference in the shoot growth among the five varieties. Furthermore, shoot production increased until 18 days, and declined there after (Fig. 2).

TABLE 3

THREE-WAY INTERACTION AMONGS VARIETY, SOIL MEDIA AND HORMONE FOR NUMBER OF SHOOTS AT 9 DAP OF CUTTING OF MALAY APPLE

Variety	Hormone			
	IAA 1000 ppm		NAA 1000 ppm	
	Top soil+ Sand	Sawdust + Sand	Top soil+ Sand	Sawdust + Sand
Malay Apple Jati	1.2±0.4 (1.00)	0.7±0.0 (0)	0.7±0.0 (0)	0.7±0.0 (0)
Malay Apple Bottle	0.7±0.0 (0)	1.6±0.8 (2.33)	1.4±0.6 (1.67)	0.7±0.0 (0)
Malay Apple Bagea	1.9±0.3 (3.33)	1.4±0.6 (1.67)	2.0±0.3 (3.67)	1.9±0.3 (3.33)
Malay Apple Dark Purple	1.4±0.6 (1.67)	1±0.5 (0.67)	0.7±0.0 (0)	1±0.5 (0.67)
Malay Apple Red Circle	1.1±0.7 (1.00)	1.1±0.3 (0.67)	1.2±0.4 (1.00)	0.7±0.0 (0)

Values are transformed to square root values; values within the parentheses are actual means.

TABLE 4

THREE-WAY INTERACTION AMONGS VARIETY, SOIL MEDIA AND HORMONE FOR NUMBER OF SHOOTS AT 12 DAP OF CUTTING OF MALAY APPLE

Variety	Hormone			
	IAA 1000 ppm		NAA 1000 ppm	
	Top soil+ Sand	Sawdust + Sand	Top soil+ Sand	Sawdust + Sand
Malay Apple Jati	2.2±0.5 (4.33)	1±0.5 (0.67)	1.2±0.4 (1)	1.8±1.1 (3.67)
Malay Apple Bottle	2.0±0.7 (4)	2.5±0.4 (5.67)	2.6±0.3 (6.33)	1.6±0.8 (2.33)
Malay Apple Bagea	2.5±0.3 (5.67)	2±1.3 (4.67)	2.9±0.1 (7.67)	2.8±0.1 (7.33)
Malay Apple Dark Purple	2.2±0.3 (4.33)	2.2±0.1 (4.33)	2.3±0.2 (5)	2.1±0.2 (4)
Malay Apple Red Circle	1.6±0.7 (2.33)	1.8±0.3 (2.67)	1.9±0.7 (3.33)	1.7±0.9 (3)

Values are transformed to sqrt values; value within the parenthesis is the actual mean.




Fig. 2. Number of shoots produced by malay apple varieties

### C. Root Volume

The volume of roots produced by cuttings was significantly influenced only by the interaction between potting media and hormone, but not by single factor alone (Table 5). The results showed that NAA combined with sawdust + sand mixture had the highest volume of roots and was significantly greater than other combinations of the factors, except with IAA and top soil + sand mixture (Fig. 3).

TABLE 5  
SOURCE OF VARIATION, DEGREES OF FREEDOM AND MEAN SQUARES TAKEN FROM ANALYSIS OF VARIANCE (ANOVA) FOR ROOT VOLUME IN THE LAST OF EXPERIMENT

Source of variation	df	Root volume
v	4	0.3884
m	1	1.3222
h	1	0.4242
v*m	4	0.0965
v*h	4	0.2644
m*h	1	2.7524*
v*m*h	4	0.6096
R	2	0.6327
Error	38	0.4367
CV		47.28

1/ v – variety; m – soil media; and h – rooting hormone;

\* : significant at p=0.05; \*\* : significant at p=0.01; \*\*\* : significant at p=0.001




Fig. 3. Root volume of malay apple cuttings as influenced by the two-way interaction between potting media and hormone.

The propagation of malay apple is considered as less successful by both cutting and budding methods [22]. However, other species of the Myrtaceae family have been subjected to propagation studies in Australia and some European countries like Belgium and Germany with *Syzygium paniculatum* Gaertn, a species native to Australia by stem cuttings as the ornamental plants [7]. In the current study, the cuttings of different varieties of malay apple responded differently to treatments. The five varieties of malay apple took short period to produce cuttings: most effective to use malay apple Bagea. Typically, this variety of malay apple needs more attention when compared to other varieties, especially with rot caused by fungal attacks on the cuttings [15]. In contrast, Jati was variety showed very slow appearance the first shoot, but in the last, this variety can produce better roots compared to other varieties of malay apple. The performance of malay apple variety Red circle had the lowest number of shoots in the beginning, but gave good rooting at last.

The differences among varieties can be attributed to different mechanisms to adaptation [14], which include the rate of metabolism process during producing shoots and roots ([8], [9], [15],[21], [24]). The condition of carbohydrate status ([13], [15]) in the cuttings may also influence the root and shoot production and performance of malay apple. It has been observed that leaves that already

exist on cuttings fall during root inducing period, and this could be assumed as a mechanism to allocate more energy for root initiation [15].

Although there were insignificant effects of the main factors, potting media composed of saw dust and sand media with 1:1 proportion promoted shooting and rooting. Even top soil + sand induced shoots more than seen as an interaction with hormone (NAA 1000 ppm) ([15], [25]). This was probably due to available nutrients ([18], [22], [27]) to cuttings. In contrast, the organic matter [15] in saw dust need time to decompose before ready to be used by plants. Rooting is usually influenced by physical properties of soil ([14],[15]), and hence found saw dust has a better influence than top soil due to its pore spaces.

With all observations during the experiment, the influence of hormone on shooting of cuttings was relatively similar, although NAA 1000 ppm influenced number of shoots. This probably indicated the benefit of using rooting hormone to regulate propagation of mala apple cutting ([15] [25]).

The results of this study found an interaction among variety, soil media and rooting hormone on the time to appearance of the first shoot and number of shoot at 9 and 12 days after planting. This indicates that such factors combined to stimulate rooting and shooting of malay apple cuttings. It was seen that exogenous hormone do not rapidly influence the growing of cutting, because the activity of exogenous hormone will depend on the metabolic processes inside plant cell [15].

Interaction among the factors showed that the variety Bagea, soil media top soil+ sand and hormone, NAA 1000 ppm had the best combination to stimulate shoots at 9 and 12 days after planting. This indicates that malay apple variety Bagea to be more superior and supply of NAA 1000 and top soil+ sand media could stimulate the plant to produce shoots ([12], [15]).

The top soil was found to be the best for rooting media because the physical and chemical characteristics of this component consisting a combination of minerals, decayed plant matter and nutrients from previous planting enhance absorption of nutrient by plants. This is different with saw dust as it needs more time to release nutrients following microbial decomposition. On the other hand, compactness of top soil compared to saw dust not only give the cuttings standing ability but also reduce oxidation of the hormone and remain active longer [15].

Root volume was highest with the combination of saw dust + sand media and hormone NAA 1000 ppm. Insignificant difference in dependent variables between IAA and NAA indicates the potential of using such hormones for propagation of malay apple by cuttings. Yet the use of NAA appeared better than IAA. Since synthetic auxins are not destroyed by oxidases and could persist in the plant much longer, NAA appears better than IAA [12]. The results of this research were similar to findings of Lebran et al. (1998) [7] where both IAA and NAA were used for stem cutting of *Syzygium paniculatum* Gaertn., and found non-significant difference between the two hormones. However, 2000 ppm of hormones caused the highest percentage of loss of cuttings and hence the lowest rooting percentage (52%).

Selection of cuttings from suitable branches [9] such as from good fruit quality and sweetness, plant vigor, absence

of diseases [15] is important for successful propagation by cuttings [15]. In addition, the timing of collecting cuttings from parent plant [21] and storing cuttings in polyethylene bags to prevent moisture loss helps avoid failures [15].

Moreover, preparation of hormones for dipping the cuttings has been suggested as time to absorb exogenous hormone differs among types of cuttings used ([12], [15]). In the current study, cuttings were dipped in the hormone solution for 24 hours.

The difficulty of propagating malay apple by cuttings is assumed due to presence of tanniferous cells in the Myrtaceae family and calcium oxalate in the cell vacuoles [7], which could inhibit root formation in cuttings [13]. Growth of malay apple seedling has been found better under the mother trees with its shade, and therefore provision of such environment appeared better for propagation of malay apple [27]. Supply of potting media comprised of sand would enhance rooting [28] as it avoids the buildup of excessive moisture and improves aeration.

#### IV. CONCLUSIONS

Based on the results of this study, it is found that the hardwood cutting of Bagea variety with top soil+ sand mixture and NAA 1000 ppm had the best performance in root and shoot production and volume of roots for propagation of malay apple that would help successful establishment. As a replacement soil media, sawdust + sand mixture could also be used. As malay apple in Ternate islands faces extinction due to environmental changes, vegetative propagation using the above combination and distribution of plants among the farmers are recommended.

#### REFERENCES

- [1] The Wikipedia website. (2010) [Online]. Jumlah Pulau di Indonesia (Number of Islands in Indonesia). Available: [http://id.wikipedia.org/wiki/Jumlah\\_pulau\\_di\\_Indonesia](http://id.wikipedia.org/wiki/Jumlah_pulau_di_Indonesia).
- [2] The Wikipedia website. (2011a) [Online]. Indonesia. Available: <http://en.wikipedia.org/wiki/Indonesia>.
- [3] The Wikipedia website (2011b). [Online]. Alfred Russel Wallace. Available: [http://en.wikipedia.org/wiki/Alfred\\_Russel\\_Wallace](http://en.wikipedia.org/wiki/Alfred_Russel_Wallace).
- [4] The Wikipedia website. (2011c) [Online]. Georg Eberhard Rumphius. Available: [http://en.wikipedia.org/wiki/Georg\\_Eberhard\\_Rumphius](http://en.wikipedia.org/wiki/Georg_Eberhard_Rumphius).
- [5] The Wikipedia website. (2011d) [Online]. Syzygium malaccense. Available: [http://en.wikipedia.org/wiki/Syzygium\\_malaccense](http://en.wikipedia.org/wiki/Syzygium_malaccense).
- [6] The Wikipedia website. (2011e) [Online]. Nama Gunung, Tinggi dan Letaknya di Indonesia Diurutkan Berdasarkan Tinggi (The name of Mount, high and it lies in Indonesia sorted by high). Available: [http://organisasi.org/nama\\_gunung\\_tinggi\\_dan\\_letaknyadi\\_Indonesia\\_diurutkan\\_berdasarkan\\_tinggi](http://organisasi.org/nama_gunung_tinggi_dan_letaknyadi_Indonesia_diurutkan_berdasarkan_tinggi).
- [7] A. Lebrun, A.N. Toussaint, and J. Roggemans. (1998). "Description of *Syzygium paniculatum* Gaertn. 'Verlaine' and its propagation by stem cuttings," *Scientia Horticulturae*, vol. 75, pp. 103-111.
- [8] A. Toogood, *Plant Propagation*. (1999). New York, USA: American Horticultural Society. DK Publishing, Inc.
- [9] A.H. Wiese, J.A. Zalesny, D.M. Donner, and R.S.Zalesny. (2006). "Bud Removal Affects Shoot, Root, and Callus Development of Hardwood Populus Cuttings," *Silvae Genetica*, vol. 55, no. 3, pp. 141-147.
- [10] C.W. Smith. (2005). "Plant Propagation, Adapted from Texas Master gardener Manual," New Mexico State University (NMSU) Extension Horticulture Specialist.
- [11] E. Mirmanto. (2010). "Flora composition and natural forest in Ternate island, North Maluku," *Indonesia Biology Journal*, vol.6, no. 3, pp. 341-351.
- [12] F. B. Salisbury and C.W. Ross (1992). *Plant Physiology*, 4th ed., California, USA: Wadsworth Publishing Company.
- [13] H. Chiun Wu (2006). "Improving in vitro propagation of *Protea cynnaroides* L. (King Protea) and the roles of starch and phenolic compounds in the rooting of cuttings," Departement of Plant Production and Soil Science, Faculty of Natural and Agricultural Sciences, University of Pretoria, Pretoria.
- [14] H. Lambers, F. S. Chapin III, and T. L. Pons. (2008). *Plant Physiological Ecology*, 2nd ed. New York, United States: Springer Science&Business Media, LLC.
- [15] H.T. Hartmann, D.E. Kester, F.T. Davies, Jr., R. L. Geneve (2002). *Plant Propagation, Principles and Practices*. 6th ed., Prentice Hall of India. New Delhi. India.
- [16] J.E. Bidlack, and S.H. Jansky (2011). *Stern's Introductory Plant Biology*. 12th ed. United States, McGraw-Hill Companies, Inc.
- [17] J.F. Morton, (1987). "Malay Apple," in *Fruits of Warm Climates*, Miami, Florida, USA, pp. 378-381.
- [18] J.F. Redhead and J.B. Hall (1992). *Tropical Forestry*, ser. Intermediate Tropical Agriculture. Kuala Lumpur, Malaysia: Longman Group.
- [19] J.G. Harris and M.W. Harris (2001). *Plant Identification Terminology, a Illustrated Glossary*. 2nd ed. Utah, United States: Spring Lake Publishing.
- [20] K.A. Monk, Y. Fretes, and L.G. Reksodihardjo (2000). *Ecology of Nusa Tenggara and Moluccas*. Ecology Series of Indonesia, 5th Book. (Indonesian Edition under Indonesian Language Team editor Kartikasari, S.N.). Jakarta, Indonesia: Dalhousie University/CIDA-Prenhallindo.
- [21] L. Schmidt (1993). *Vegetative Propagation. Guidelines on Grafting, Air-Layering and Cuttings*. FAO. Available at: <http://www.fao.org/docrep/006/ad224e/AD224E21.htm>.
- [22] N.A. Prastowo, J.M. Rosetko, G.E.S. Maurung, E. Nugraha, J.M. Tukan, F. Harum (2006). *Teknik Pembibitan dan Perbanyakan Vegetatif Tanaman Buah*. Bogor, Indonesia: World Agroforestry Centre (ICRAF) & Winrock International.
- [23] P. M. Taylor and M. N. Richards (1999). "Ternate, The Residency and its Sultanate" translated from the Dutch "Bijdragen tot de kennis der Residentie Ternate," by F.S.A. de Clercq (1890). Washington, D.C, Smithsonian Institution Libraries Digital Edition, Available: [http://www.sil.si.edu/DigitalCollections/Anthropology/Ternate/terna te.pdf](http://www.sil.si.edu/DigitalCollections/Anthropology/Ternate/terna_te.pdf).
- [24] R. Leatherwood, A. Enfield, and R. Lopez. (2006). *Crop Cultivation, Getting the Most Out of Cuttings*. North Carolina State and Michigan State Universities, GPN, United States.
- [25] R.K. Tiwari, and K. Das (2010). "Effect of stem cutting and hormonal pre-treatment on propagation of *Embelia tsjeriam* and *Caesalpinia bonduc*, two important medical plant species," *Journal of Medicinal Plants Research*, vol. 4, no. 15, pp. 1577-1583.
- [26] R.M. Crasweller (2005). *Grafting and Propagating Fruit Trees*. United States, The Pennsylvania State University, College of Agricultural Sciences.
- [27] V. Kumar (1999). *Nursery and Plantation Practices in Forestry*. 2nd ed. Jodhpur, India: Scientific Publishers.
- [28] W.A. Whistler, and C.R. Elevitch (2006). *Syzygium malaccense* (Malay apple), ver. 2.1. In: Elevitch, C.R. (ed.). *Species Profiles for Pacific Island Agroforestry*. Permanent Agriculture Resources (PAR), Holualoa, Hawaii. Available at: <http://www.agroforestry.net/tti/Syzygium-Malayapple.pdf>.