













## REFERENCES

- [1] K.K. Tripathi, R. Warriar, O.P. Govila, V. Ahuja, *Biology of Zea mays (Maize)*, Dep. Biotechnol. Ministry of Science and Technology & Ministry of Environment and Forests, Gov. India, 2011.
- [2] P. Ranum, J.P. Peña -Rosas, and M.N. Garcia-Casal, "Global maize production, utilization, and consumption," *Ann.N.Y.Acad.Sci.*, vol. 1312(1), pp.105-112, 2014.
- [3] Statistic Indonesia, "Foreign Trade Statistical Bulletin:Imports," Jakarta, Indonesia; BPS Statistic Indonesia, 2014.
- [4] Statistic Indonesia, "Foreign Trade Statistical Bulletin:Impors," Jakarta, Indonesia; BPS Statistic Indonesia, 2015.
- [5] K.H. Tan, *Soils in the Humid Tropics and Monsoon Region of Indonesia*, Florida, USA: CRC Press, 2008.
- [6] N. Hakim, R. Alfina, H. Agustian, and Yulnafatmawita, "Bacterial inoculants to increase the biomass and nutrient uptake of tithonia cultivated as hedgerow plants in Ultisols," *Malays J. Soil Sci.*, vol. 18, pp. 115-123, 2014.
- [7] E.C. Omondi, J.B. Norton and D.S. Ashilenje, "Performance of a local open pollinated maize variety and a common hybrid variety under intensive small-scale farming practices," *Afr. J. Agric. Res.*, vol.9 (11), pp. 950-955, 2014.
- [8] Y. Soelaeman and U. Haryati, "Soil physical properties and production of upland Ultisol soil as influenced by manure application and P fertilization," *Agrivita*, vol. 34(2), pp. 136-143, 2012.
- [9] M. Syukur, S. Sujiprihati, R. Yunianti, dan D. A. Kusumah, "Evaluasi daya hasil cabai hibrida dan daya adaptasinya di empat lokasi dalam dua tahun," *J. Agron. Indonesia*, vol. 38(1), pp. 43-51, 2010.
- [10] H. G. Gauch, "Statistical analysis of yield trials by AMMI and GGE," *Crop Sci*, vol. 46(4), pp. 1488-1500, 2006.
- [11] P. H. C. de Mattos, R. A. de Oliveira, J. C. B. Filho, E. Daros and M. A. A. Veríssimo, "Evaluation of sugarcane genotypes and production environments in Paraná by GGE biplot and AMMI analysis," *Crop Breed. and Appl. Biotechnol.*, vol.13(1), pp. 83-90, 2013.
- [12] R. W. Zobel, M. J. Wright and H. G. Gauch, "Statistical analysis of a yield trial," *Agron. J.*, vol. 80(3), pp. 388-393, 1988.
- [13] P.S. Rao, P.S. Reddy, A. Rathore, B.V. Reddy and S. Panwar, "Application GGE biplot and AMMI model to evaluate sweet sorghum (*Sorghum bicolor*) hybrids for genotype x environment interaction and seasonal adaptation," *Indian J. Agric. Sci.*, vol. 81(5), pp. 438-444, 2011.
- [14] A. Khalil, H. Rahman, N. U. Rehman, M. Arif, I. H. Khalil, M. Iqbal, Hidayatullah, K. Afridi, M. Sajjad, and M. Ishaq, "Evaluation of maize hybrids for grain yield stability in north-west of Pakistan," *Sarhad J. Agric.*, vol. 27(2), pp. 213-218, 2011.
- [15] M. Munawar, G. Hammad and M. Shahbaz, "Evaluation of maize (*Zea mays* L.) hybrids under different environments by GGE biplot analysis," *Am-Eurasian J. Agric. Environ. Sci.*, vol. 13(9), pp. 1252-1257, 2013.
- [16] Rustikawati, E. Supriyono, A. Romeida, C. Herison, S.H. Sutjahjo, "Identification of M4 gamma irradiated maize mutant based on RAPD markers" *Agrivita*, vol.34(2), pp. 161-165, 2012.
- [17] B. Setyawan, I. Suliansyah, A. Anwar, E. Swasti, "Preliminary trial of 11 new hybrid maize genotype to the resistance on Java Downy Mildew (*Peronosclerospora maydis*)," *Int. J. Adv. Sci. Eng. Inf. Technol.* vol. 6(2), pp. 262-264, 2016.
- [18] B. Setyawan, I. Suliansyah, A. Anwar, E. Swasti, "Agronomic characters, yield components and grain yield evaluation of 11 new hybrid maize prospective genotypes," *Int. J. Adv. Sci. Eng. Inf. Technol.* vol. 6(4), pp. 483-488, 2016.
- [19] H. G. Gauch, H. P. Piepho, and P. Annicchiarico, "Statistical analysis of yield trials by AMMI and GGE: Further considerations," *Crop Sci*, vol. 48(3), pp. 866-889, 2008.
- [20] M. Vargas, J. Crossa, K. Sayre, M. Reynolds, M. E. Ramirez, M. Talbot, "Interpreting genotype x environment interaction in wheat by Partial Least Square Regression," *Crop Sci*. vol. 38(3), pp. 679-689, 1998.
- [21] J. N. Nwite, "Predicting grain yields of maize in an Ultisol amended with organic wastes using modified productivity index in Abakaliki, Southeastern Nigeria," *Afr. J. Agric. Res.*, vol 11(44), pp. 4434-4443, 2016.
- [22] A.O. Akongwubel, U. B. Ewa, A. Prince, O. Jude, A. Martins, O. Simon, O. Nicholas, "Evaluation of agronomic performance of maize (*Zea mays* L.) under different rates of poultry manure application in an Ultisol of Obubra, Cross River State, Nigeria," *Int. J. Agric. For.*, vol 2(4), pp.138-144, 2012.
- [23] C. Herison, M. Handajningsih, Fahrurrozi, and Rustikawati, "Wet season trials on growth and yield of six newly developed chili pepper hybrids at three different locations," *Int. J. Adv. Sci. Eng. Inf. Technol.* vol. 7(5), pp. 913-919, 2017.
- [24] A. A. Mattjik, I. M. Sumertajaya, A.F. Hadi, and G. N. A. Wibawa. Additive Main-effect and Multiplicative Interaction (AMMI) Modeling : Present and Future (in Indonesian), IPB Press, 2011.
- [25] B. Mitrović, D. Stanisavljevi, S. Treski, M. Stojaković, M. Ivanović, G. Bekavac, M. Rajković, "Evaluation of experimental maize hybrids tested in multi-location trials using AMMI and GGE biplot analyses," *Turk. J. Field Crops*, vol. 17(1), pp. 35-40, 2012.
- [26] E. T. Nuss, S. A. Tanumihardjo, "Maize: a paramount staple crop in the context of global nutrition" *Compr. Rev. in Food Sci. Food Saf.* vol. 9(4), pp.417-436, 2010.