

- [11] J. V. Holst, G. F. Versteeg, D. W. F. Brillman, and J. A. Hogendoorn, "Kinetic study of CO₂ with various amino acid salts in aqueous solution," *Chem. Eng. Sci.*, vol. 64, pp. 59–68, 2009.
- [12] H. Thee, K. H. Smith, G. da Silva, S. E. Kentish, and G. W. Stevens, "Carbon dioxide absorption into unpromoted and borate-catalyzed potassium carbonate solutions," *Chem. Eng. J.*, vol. 181–182, pp. 694–701, 2012.
- [13] S. Shen, X. Feng, R. Zhao, U. K. Ghosh, and A. Chen, "Kinetic study of carbon dioxide absorption with aqueous potassium carbonate promoted by arginine," *Chem. Eng. J.*, vol. 222, pp. 478–487, 2013.
- [14] B. Prasad, "Experimental investigation of CO₂ absorption capacity in amine blend of 2-Amino-2-methyl-1-propanol (AMP) and Tetraethylenepentamine (TEPA)," Master thesis, Thapar University, India, 2013.
- [15] S. Paul and K. Thomsen, "Kinetics of absorption of carbon dioxide into aqueous potassium salt of proline," *Int. J. Greenh. Gas Control*, vol. 8, pp. 169–179, 2012.
- [16] M. Gupta, F. Eirik, and H. F. Svendsen, "Computational study of equilibrium constants for amines and amino acids for CO₂ capture solvents," *Energy Procedia*, vol. 37, pp. 1720–1727, 2013.
- [17] C. A. Fitch, G. Platzer, M. Okon, B. G. E, and L. P. McIntosh, "Arginine: Its pKa value revisited," *Proteine Sci.*, vol. 24, no. 5, pp. 752–761, 2015.
- [18] K. A. Hoff, "Modeling and experimental study of carbon dioxide absorption in a membrane contactor," Doctoral thesis, Norwegian University of Science and Technology, Norway, 2003.
- [19] T. J. Edwards, G. Maurer, J. Newman, and J. M. Prausnitz, "Vapor-liquid equilibria in multicomponent aqueous solutions of volatile weak electrolytes," *AIChE J.*, vol. 24, no. 6, pp. 900–976, 1978.
- [20] G. Xu, C. Zhang, S. Qin, W. Gao, and H. Liu, "Gas-liquid equilibrium in a CO₂-MDEA-H₂O system and the effect of piperazine on it," *Ind. Eng. Chem. Res.*, vol. 37, pp. 1473–1477, 1998.
- [21] D. Barth, C. Tondre, and J. J. Delpuech, "Kinetics and mechanism of the reactions of carbon dioxide with alkanolamines: A discussion concerning the cases of MDEA and DEA," *Chem. Eng. Sci.*, vol. 39, no. 12, pp. 1753–1757, 1984.
- [22] P. V. Danckwerts, *Gas liquid reactions*. New York: McGraw-Hill, 1970.
- [23] C. J. Geankoplis, *Transport processes and unit operations*, 3rd editio. New Delhi: Prentice-Hall of India, 1997.
- [24] J. T. Cullinane and G. T. Rochelle, "Carbon dioxide absorption with aqueous potassium carbonate promoted by piperazine," *Chem. Eng. Sci.*, vol. 59, pp. 3619–3630, 2004.
- [25] F. Yi, H.-K. Zou, G.-W. Chu, L. Shao, and J.-F. Chen, "Modeling and experimental studies on absorption of CO₂ by Benfield solution in rotating packed bed," *Chem. Eng. J.*, vol. 145, no. 3, pp. 377–384, 2009.
- [26] C. R. Sander, "Compilation of Henry's law constants for inorganic and organic species of potential importance in environmental chemistry," *Max Planck Inst. Chem. J.*, vol. 3, p. 57, 1999.
- [27] H. Kierzkowska-pawlak, M. Siemieniec, and A. Chacuk, "Reaction kinetics of CO₂ in aqueous methyldiethanolamine solutions using the stopped-flow technique," *Chem. Process Eng.*, vol. 33, no. 1, pp. 7–18, 2011.
- [28] M. P. Gimeno, M. C. Mayoral, and J. M. Andrés, "Influence of temperature on CO₂ absorption rate and capacity in ionic liquids," *Energy & Fuels*, vol. 27, no. 7, pp. 3928–3935, 2013.
- [29] Z. Feng, F. Cheng-Gang, W. You-Ting, W. Yuan-Tao, L. Ai-Min, and Z. Zhi-Bang, "Absorption of CO₂ in the aqueous solutions of functionalized ionic liquids and MDEA," *Chem. Eng. J.*, vol. 160, no. 2, pp. 691–697, 2010.
- [30] J. A. Bullin, J. C. Polasek, and S. T. Donnelly, "The use of MDEA and mixtures of amines for bulk CO₂ removal," in *Proceedings of the Sixty-Ninth GPA Annual Convention*, 1990, pp. 1–9.
- [31] N. Ramachandran, A. Aboudheir, R. Idem, and P. Tontiwachwuthikul, "Kinetics of the absorption of CO₂ into mixed aqueous loaded solutions of monoethanolamine and methyldiethanolamine," *Ind. Eng. Chem. Res.*, vol. 45, no. 8, pp. 2608–2616, 2006.
- [32] M. Edali, R. Idem, and A. Aboudheir, "1D and 2D absorption-rate/kinetic modeling and simulation of carbon dioxide absorption into mixed aqueous solutions of MDEA and PZ in a laminar jet apparatus," *Int. J. Greenh. Gas Control*, vol. 4, pp. 143–151, 2010.